



# Need more money for college expenses?

The CLC Private Loan<sup>SM</sup> can get you up to  
**\$40,000** a year for college-related expenses.

---

Here's why the CLC Private Loan<sup>SM</sup> is a smart choice:

- ✓ Approved borrowers are sent a check within four business days
- ✓ Get \$1000 - \$40,000 each year
- ✓ Interest rates as low as prime + 0% (8.66% APR)
- ✓ Quick and easy approval process
- ✓ No payments until after graduating or leaving school

**CLICK HERE**

or call **800.311.9615.**

*We are available 24 hours  
a day, 7 days a week.*

# Polynomials and Exponents

If Diane makes \$13 per hour and works 40 hours per week, she will earn more than one million dollars over the next 40 years. How much money will Diane earn? Perform the computations using scientific notation.



## 3-1 ■ Exponents—I

### Exponential form

In chapter 1, we discussed the idea of exponents as related to real numbers. Since variables are symbols for real numbers, let us now apply the idea of exponents to variables. The expression  $x^4$  is called the **exponential form** of the product

$$x \cdot x \cdot x \cdot x$$

We call  $x$  the **base** and 4 the **exponent**.

$$\begin{array}{c}
 \text{Exponent} \\
 \downarrow \\
 \text{Exponential form} \rightarrow x^4 = x \cdot x \cdot x \cdot x \leftarrow \text{Expanded form} \\
 \uparrow \qquad \qquad \qquad \downarrow \\
 \text{Base} \qquad \qquad \qquad \text{4 factors of } x
 \end{array}$$

### Definition of exponents

$$a^n = \underbrace{a \cdot a \cdot a \cdots a}_{n \text{ factors of } a}, \text{ where } n \text{ is a positive integer.}$$

### Concept

The exponent tells us how many times the base is used as a factor in an indicated product.

**Note** An exponent acts only on the symbol immediately to its left. That is, in  $ab^4$  the exponent 4 applies only to  $b$ , whereas  $(ab)^4$  means the exponent applies to both  $a$  and  $b$ .



**Example 3-1 A**

Write in exponential form.

1.  $2 \cdot 2 \cdot 2 \cdot 2 = 2^4$

2.  $a \cdot a \cdot a = a^3$

3.  $(a + b)(a + b)(a + b) = (a + b)^3$

**Note** In example 3,  $(a + b)$  is the base.

4.  $(-3)(-3)(-3)(-3) = (-3)^4$

5.  $-(3 \cdot 3 \cdot 3 \cdot 3) = -3^4$

**Note** Examples 4 and 5 review the ideas from section 1-6 on exponents related to real numbers. Recall that  $(-3)^4 = 81$ , whereas  $-3^4 = -81$ .**Quick check** Write  $y \cdot y \cdot y \cdot y$  in exponential form. ■**Example 3-1 B**

Write as an indicated product.

1.  $b^4 = b \cdot b \cdot b \cdot b$

2.  $5^3 = 5 \cdot 5 \cdot 5$

3.  $(x - y)^4 = (x - y)(x - y)(x - y)(x - y)$

4.  $(-2)^2 = (-2)(-2)$

5.  $-2^2 = -(2 \cdot 2)$

**Quick check** Write  $c^5$  as an indicated product. ■**Multiplication of like bases**Consider the indicated product of  $x^2 \cdot x^3$ . If we rewrite  $x^2$  and  $x^3$  by using the definition of exponents, we have

$$x^2 \cdot x^3 = \overbrace{x \cdot x}^{x^2} \cdot \overbrace{x \cdot x \cdot x}^{x^3}$$

and again using the definition of exponents, this becomes

$$x^2 \cdot x^3 = \overbrace{x \cdot x \cdot x \cdot x \cdot x}^{5 \text{ factors}} = x^5$$

This leads us to the observation that

$$\begin{array}{c}
 \text{Add exponents} \\
 \downarrow \\
 x^2 \cdot x^3 = x^{2+3} = x^5 \\
 \begin{array}{cc}
 \uparrow & \uparrow \\
 \text{Multiply} & \text{Base remains} \\
 \text{like bases} & \text{unchanged}
 \end{array}
 \end{array}$$

Thus we have the following **product property of exponents**.**Product property of exponents**

$$a^m \cdot a^n = a^{m+n}$$

**Concept**

When multiplying like bases, add their exponents.

**Note** The base stays the same throughout the process. It is by adding the exponents that the multiplication is carried out.

### Example 3-1 C

Find the product.

$$1. x^3 \cdot x^5 = x^{3+5} = x^8$$

$$2. 3^2 \cdot 3^4 = 3^{2+4} = 3^6 = 729$$

**Note** A common error in multiplying  $3^2 \cdot 3^4$  is to multiply the bases  $3 \cdot 3 = 9$  and add the exponents, getting the incorrect answer of  $9^6$ . The correct way is to say  $3^2 \cdot 3^4 = 3^6$ , not  $9^6$ .

$$3. y^2 \cdot y^3 \cdot y^4 = y^{2+3+4} = y^9$$

$$4. a^2 \cdot a \cdot a^3 = a^{2+1+3} = a^6$$

**Note** The variable  $a$  means the same as  $a^1$ . Likewise, 3 means the same as  $3^1$ . If there is no exponent written with a numeral or a variable, the exponent is understood to be 1.

$$5. (a + b)^3(a + b)^4 = (a + b)^{3+4} = (a + b)^7$$

$$6. (-2)^3(-2)^2 = (-2)^{3+2} = (-2)^5 = -32$$

► **Quick check** Find the product.  $x^4 \cdot x^5$

### Group of factors to a power property of exponents

Several additional properties of exponents can be derived using the definition of exponents and the commutative and associative properties of multiplication. Observe the following:

$$\begin{aligned} (xy)^3 &= \overbrace{xy \cdot xy \cdot xy}^{3 \text{ factors of } xy} \\ &= \overbrace{x \cdot x \cdot x}^{3 \text{ factors of } x} \cdot \overbrace{y \cdot y \cdot y}^{3 \text{ factors of } y} \\ &= x^3 y^3 \end{aligned}$$

This leads us to the following property of exponents.

### Group of factors to a power property of exponents

$$(ab)^n = a^n b^n$$

#### Concept

When a group of *factors* is raised to a power, raise each of the factors in the group to this power.

### Example 3-1 D

Simplify.

$$1. (ab)^4 = a^4 b^4$$

Both  $a$  and  $b$  are raised to the 4th power

$$2. \quad \begin{array}{ccccccc} \text{Groups of factors to a power} & & \text{Raise each factor to the power} & & \text{Standard form} \\ (2ab)^3 & = & 2^3 a^3 b^3 & = & 8a^3 b^3 \end{array}$$

**Note** In example 2, the number 2 is a factor in the group. Therefore it is also raised to the indicated power.



$$3. (3 \cdot 4)^3 = 3^3 \cdot 4^3 = 27 \cdot 64 = 1,728 \quad (3 \cdot 4)^3 \text{ also is } (12)^3 = 1,728$$

**Note** The quantity  $(a + b)^3 \neq a^3 + b^3$  because  $a$  and  $b$  are *terms*, not factors as the property specified. If we consider  $(a + b)$  to be a single factor, then by the definition of exponents we have

$$(a + b)^3 = (a + b)(a + b)(a + b)$$

We will see the method of multiplying this later in this chapter.

### Power of a power

Consider the expression  $(x^4)^3$ . Applying the definition of exponents and the product property of exponents, we have

$$(x^4)^3 = \overbrace{x^4 \cdot x^4 \cdot x^4}^{3 \text{ factors of } x^4} = \overbrace{x^{4+4+4}}^{\text{Add the exponents}} = x^{12}$$

In chapter 1, we reviewed the idea that multiplication is repeated addition of the same number. Therefore adding the exponent 4 three times is the same as  $4 \cdot 3$ . Thus

$$(x^4)^3 = \overbrace{x^4 \cdot x^4 \cdot x^4}^{\text{Power of a power}} = \overbrace{x^4 \cdot 3}^{\text{Multiply exponents}} = x^{12}$$

Therefore we have the following property of exponents.

### Power of a power property of exponents

$$(a^m)^n = a^{m \cdot n}$$

#### Concept

A power of a power is found by multiplying the exponents.

### Example 3-1 E

Simplify.

$$1. (y^3)^2 = y^3 \cdot 2 = y^6$$

$$2. (4^2)^5 = 4^2 \cdot 5 = 4^{10} = 1,048,576$$

$$3. (x^5)^4 = x^5 \cdot 4 = x^{20}$$

► **Quick check** Simplify.  $(a^4)^3$

### Products of monomials

To multiply the monomials

$$3x^2 \cdot 5x$$

we apply the commutative and associative properties of multiplication along with the properties of exponents. We then write this expression as a product of the numerical coefficients times the product of the variables. That is,

$$3x^2 \cdot 5x = (3 \cdot 5)(x^2 \cdot x) = 15x^3$$

To find the product of

$$5a \cdot 4b$$

we apply the same properties to get

$$5a \cdot 4b = (5 \cdot 4)(a \cdot b) = 20ab$$

**Note** It is a good procedure to write the variable factors of any term in alphabetical order. This makes identifying like terms much simpler. For example,  $3a^2c^3b$  and  $4bc^3a^2$  are like terms, but recognizing that fact would have been easier if they had been written as  $3a^2bc^3$  and  $4a^2bc^3$ .

### ■ Example 3-1 F

Perform the indicated multiplication.

- $4x \cdot 3xy = (4 \cdot 3) \cdot (x \cdot x) \cdot y = 12x^2y$
- $8a^3 \cdot 4a^3 \cdot 3a = (8 \cdot 4 \cdot 3) \cdot (a^3 \cdot a^3 \cdot a) = 96a^7$
- $(-2a^2) \cdot (3ab) = (-2 \cdot 3) \cdot (a^2 \cdot a) \cdot b = -6a^3b$

**Note** The product of  $a^3$  and  $b$  can *only* be written as  $a^3b$  since  $a$  and  $b$  are not like bases.

$$4. (5x^2y^3z)(4x^3yz^4) = (5 \cdot 4)(x^2x^3)(y^3y)(zz^4) = 20x^5y^4z^5$$

### Problem solving

The following problems require us to write algebraic expressions involving the use of exponents.

### ■ Example 3-1 G

Write an algebraic expression for each of the following verbal statements.

- The volume of a cube is found by using the length of the edge,  $e$ , as a factor 3 times. Write an expression for the volume of a cube.

We write  $e$  as a factor 3 times as  $e \cdot e \cdot e = e^3$ . Then the volume,  $V$ , of a cube is given by

$$V = e^3.$$

- Write an expression for 5 less than the square of a number.

Let  $n$  represent the number, then the square of the number is given as  $n^2$ , and since “less than” means to subtract, the expression is given by

$$n^2 - 5.$$

### Mastery points

Can you

- Write a product in exponential form?
- Use the product property of exponents?
- Raise a group of factors to a power?
- Raise a power to a power?
- Multiply monomials?



**Exercise 3-1**

Write the following expressions in exponential form. See example 3-1 A.

**Example**  $y \cdot y \cdot y \cdot y$

**Solution**  $= y^4$   $y$  to the fourth power

1.  $aaaaa$
2.  $bbbb$
3.  $(-2)(-2)(-2)(-2)$
4.  $-(2 \cdot 2 \cdot 2 \cdot 2)$
5.  $xxxxxx$
6.  $(2a)(2a)(2a)$
7.  $(xy)(xy)(xy)(xy)$
8.  $(a + b)(a + b)$
9.  $(x - y)(x - y)(x - y)$
10.  $(2a - b)(2a - b)(2a - b)$

Write as an indicated product (expanded form). See example 3-1 B.

**Example**  $c^5$

**Solution**  $= c \cdot c \cdot c \cdot c \cdot c$   $c$  written as a factor 5 times

11.  $x^4$
12.  $y^5$
13.  $(-2)^3$
14.  $-2^4$
15.  $5^3$
16.  $(5x)^3$
17.  $(4y)^4$
18.  $(a + b)^3$
19.  $(x - y)^2$
20.  $(2x + y)^3$

Simplify by using the properties of exponents. See examples 3-1 C, D, E, and F.

**Examples**  $x^4 \cdot x^5$

**Solutions**  $= x^{4+5}$   
 $= x^9$

Like bases  
Add exponents

$(a^4)^3$

$= a^{4 \cdot 3}$   
 $= a^{12}$

Power of a power  
Multiply exponents

21.  $x^4 \cdot x^7$
22.  $a^5 \cdot a^5$
23.  $R^2 \cdot R$
24.  $a \cdot a^4$
25.  $a^2 \cdot a^3 \cdot a^4$
26.  $x^5 \cdot x \cdot x^3$
27.  $5^2 \cdot 5^3$
28.  $6 \cdot 6^3$
29.  $4 \cdot 4^2 \cdot 4^4$
30.  $(a + b)^2(a + b)^5$
31.  $(x - 2y)^4(x - 2y)^6$
32.  $(3a + b)^2(3a + b)^3$
33.  $(a - b)^4(a - b)^7$
34.  $(ab)^5$
35.  $(xy)^4$
36.  $(2abc)^3$
37.  $(4xyz)^3$
38.  $(a^2)^4$
39.  $(x^5)^3$
40.  $(y^2)^2$
41.  $(b^5)^5$
42.  $(c^9)^3$
43.  $(2xy^2)(3x^3y)$
44.  $(4x^2y^3)(5xy^4)$
45.  $(a^2b^3)(a^5b^2)$
46.  $(x^2y^2)(x^4y^3)$
47.  $(6x^3)(5x^2)$
48.  $(4a)(3a^4)$
49.  $(2a^3b^4c)(6a^4b^3)$
50.  $(5xy)(xy)$
51.  $(3a^2b)(4a^3b^2)$
52.  $(a^3b^4)(5a^2b^5)$
53.  $(-2a^2b)(3ab^4)$
54.  $(-5x^2y^5)(-2x^2y)$

55. The formula for finding the volume of a cube is  $V = e^3$ , where  $V$  represents volume in some cubic unit of measure and  $e$  represents the length of the edge of the cube. Write an expression for the volume in expanded form, and then determine the number of cubic units in the figure for each of the following values of  $e$ : (a)  $e = 5$ , (b)  $e = 4$ , (c)  $e = 6$ .

# Love The Taste. Taste The Love.

At Culver's® we can't think of anything better than serving up our creamy frozen custard and delicious classics cooked fresh the minute you order them. Which is why when we bring them to your table, they're always accompanied by a warm smile and a friendly offer to see if there's anything else we can get for you. So come on into your neighborhood Culver's and see for yourself. You might just be in love by the time you leave.





Using exponents, write an expression for each of the following verbal statements. See example 3-1 G.

56. The area,  $A$ , of a square is found by using the length of the side,  $s$ , as a factor twice. Write an expression for the area of a square.
57. The distance,  $s$ , a falling object will fall in time,  $t$ , seconds is found by multiplying  $\frac{1}{2}$  times the gravity,  $g$ , times the square of  $t$ . Write an expression for the distance the object will fall.
58. The area of a circle is found by multiplying the constant  $\pi$  times the length of the radius,  $r$ , used as a factor 2 times. Write an expression for the area of a circle.
59. The volume,  $V$ , of a sphere is found by multiplying  $\frac{4}{3}\pi$  times the radius,  $r$ , used as a factor 3 times. Write an expression for the volume of a sphere.
60. Johnny is  $n$  years old. His mother says that she is 6 years more than the cube of Johnny's age. Write an expression for his mother's age.
61. Jane is  $m$  years old. Her father is 8 years less than Jane's age used as a factor 4 times. Write an expression for her father's age.
62. Write an expression for 2 times the square of  $t$ .
63. Write an expression for twice the square of  $x$  less the cube of  $y$ .
64. A number can be written in the form  $a$  times 10 used as a factor 8 times, where  $a$  is a number between 1 and 10. Write an expression for the number in terms of  $a$ .
65. Write an expression for the quotient of the cube of  $p$  divided by the square of  $q$ .

### Review exercises

Perform the indicated addition and subtraction. See section 2-3.

1.  $2a + 3a + 4a$
2.  $5x + x + 2x$
3.  $3ab - 2ab + 5ab$
4.  $9xy + 4xy - 6xy$
5.  $4a^2 + 3a^2 - 2a + 7a$
6.  $6x^2 + 3x - x^2 + 2x$
7.  $2x^2y - x^2y + 3xy^2 + 4xy^2$
8.  $5ab^2 + 3a^2b - 2ab^2 - a^2b$

## 3-2 ■ Products of algebraic expressions

### Product of a monomial and a multinomial

To multiply a monomial and a multinomial (a polynomial of more than one term), we use the distributive property. For example, to multiply

$$3x^2y(x^2 + xy + y^2)$$

we multiply each term in the trinomial by the monomial  $3x^2y$  to get

$$(3x^2y \cdot x^2) + (3x^2y \cdot xy) + (3x^2y \cdot y^2)$$

which yields

$$3x^2y(x^2 + xy + y^2) = 3x^4y + 3x^3y^2 + 3x^2y^3$$

In each indicated product, note that we multiplied like bases by using the properties of exponents. For example, in the first term,

$$3x^2y \cdot x^2 = 3 \cdot (x^2 \cdot x^2) \cdot y = 3 \cdot x^{2+2} \cdot y = 3x^4y$$

**Example 3-2 A**

Perform the indicated multiplication.

$$1. \quad 5y(2y + 3) = 5y \cdot 2y + 5y \cdot 3 \quad \begin{array}{l} \text{Distribute } 5y \text{ times each term in the parentheses} \\ \text{Multiply monomials} \end{array}$$

$$= 10y^2 + 15y$$

$$2. \quad x^3(x^2 + xy - y^2) = x^3 \cdot x^2 + x^3 \cdot xy - x^3 \cdot y^2$$

$$= x^5 + x^4y - x^3y^2$$

**Note** In example 2, when we multiplied  $x^3$  times the third term of the trinomial,  $y^2$ , the subtraction sign remained, giving  $-x^3y^2$ .

$$3. \quad -5a^3(a^2 + 2ab - b^3) = -5a^3 \cdot a^2 - 5a^3 \cdot 2ab + 5a^3 \cdot b^3$$

$$= -5a^5 - 10a^4b + 5a^3b^3$$

$$4. \quad 4x^2y(2x^3 - 3x^2y^2 + y^4) = 4x^2y \cdot 2x^3 - 4x^2y \cdot 3x^2y^2 + 4x^2y \cdot y^4$$

$$= 8x^5y - 12x^4y^3 + 4x^2y^5$$

► **Quick check** Perform the indicated multiplication.  $3ab^2(2a - 3b)$

**Product of two multinomials**

The product of two multinomials requires the use of the distributive property several times. That is, in the product

$$(x + 2y)(x + y)$$

we consider  $(x + 2y)$  a single number and apply the distributive property.

$$(x + 2y)(x + y) = (x + 2y) \cdot x + (x + 2y) \cdot y$$

We now apply the distributive property again.

$$(x + 2y) \cdot x + (x + 2y) \cdot y = x \cdot x + 2y \cdot x + x \cdot y + 2y \cdot y$$

$$= x^2 + 2xy + xy + 2y^2$$

The last step in the problem is to combine like terms, if there are any.

$$x^2 + (2xy + xy) + 2y^2 = x^2 + 3xy + 2y^2$$

Notice that in this product, each term of the first factor is multiplied by each term of the second factor. We can generalize our procedure as follows:

**Multiplying two multinomials**

When we are multiplying two multinomials, we multiply each term in the first multinomial by each term in the second multinomial. We then combine like terms.

**Example 3-2 B**

Perform the indicated multiplication and simplify.

$$1. \quad (a + 3)(a - 4) = a \cdot a - a \cdot 4 + 3 \cdot a - 3 \cdot 4 \quad \begin{array}{l} \text{Distribute multiplication} \\ \text{Multiply monomials} \\ \text{Combine like terms} \end{array}$$

$$= a^2 - 4a + 3a - 12$$

$$= a^2 - a - 12$$

**Note** We have drawn arrows to indicate the multiplication that is being carried out. This should be a convenient way for us to indicate the multiplication to be performed.



$$\begin{array}{l}
 \text{2. } (2x + 3)(5x - 2) = 10x^2 - 4x + 15x - 6 \\
 \phantom{2. } = 10x^2 + 11x - 6
 \end{array}$$

Distribute and multiply  
Combine like terms

**Note** A word that is useful for remembering the multiplication to be performed when multiplying two binomials is **FOIL**. Foil is an abbreviation signifying **F**irst times **f**irst, **O**uter times **o**uter, **I**nner times **i**nner, and **L**ast times **l**ast.

$$\begin{array}{l}
 \text{3. } (a + b)(a + 2b) = a^2 + 2ab + ab + 2b^2 \\
 \phantom{3. } = a^2 + 3ab + 2b^2
 \end{array}$$

Distribute and multiply  
Combine like terms

► **Quick check** Perform the indicated multiplication and simplify.  
 $(2x + y)(x - 3y)$

### Special products

Three special products appear so often that the form of the answers can be written without computation. Consider the product

$$(x + 6)^2 = (x + 6)(x + 6)$$

which becomes

$$x^2 + 6x + 6x + 36$$

When we combine the second and third terms, we get

$$x^2 + 12x + 36$$

This is called the **square of a binomial** or a **perfect square trinomial** and has certain characteristics. Inspection shows us that in

$$(x + 6)^2 = x^2 + 12x + 36$$

the three terms of the product can be obtained in the following manner:

#### The square of a binomial

1. The first term of the product is the *square of the first term* of the binomial  $[(x)^2 = x^2]$ .
2. The second term of the product is *two times the product of the two terms of the binomial*  $[2(x \cdot 6) = 12x]$ .
3. The third term of the product is the *square of the second term* of the binomial  $[(6)^2 = 36]$ .

If we apply this to

$$\begin{array}{l}
 (x - 7)^2 \\
 = [x + (-7)]^2 \\
 = x^2 + [2 \cdot x \cdot (-7)] + (-7)^2
 \end{array}$$

we get

and so

$$\begin{array}{l}
 (x - 7)^2 = x^2 + (-14x) + 49 \\
 \phantom{(x - 7)^2} = x^2 - 14x + 49
 \end{array}$$

In general, for real numbers  $a$  and  $b$ ,

$$(a + b)^2 = a^2 + 2ab + b^2$$

and

$$(a - b)^2 = a^2 - 2ab + b^2$$

**Note**  $(a + b)^2 = a^2 + 2ab + b^2$ , not  $a^2 + b^2$ . This is a common error. The square of a binomial is always a trinomial.

### ■ Example 3-2 C

Perform the indicated multiplication and simplify.

$$1. (2x + 3)^2 = (2x)^2 + (2 \cdot 2x \cdot 3) + (3)^2$$

$$= 4x^2 + 12x + 9$$

Apply special products  
property

Multiply monomials

$$2. (5a - 4b)^2 = (5a)^2 - [2 \cdot 5a \cdot (4b)] + (4b)^2$$

$$= 25a^2 - [40ab] + 16b^2$$

$$= 25a^2 - 40ab + 16b^2$$

Special products property

Multiply monomials

Standard form

The third special product is obtained by multiplying the sum and the difference of the same two terms. Consider the following:

$$\begin{aligned}(x + 3)(x - 3) &= x^2 - 3x + 3x - 9 \\ &= x^2 - 9\end{aligned}$$

Special characteristics are evident in this product also.

### The difference of two squares

For real numbers  $a$  and  $b$ ,

$$(a + b)(a - b) = a^2 - b^2$$

#### Concept

1. The product is obtained by first squaring the first term of the factors, and then
2. subtracting the square of the second term of the factors.

### ■ Example 3-2 D

Perform the indicated multiplication and simplify.

$$1. (x + 7)(x - 7) = (x)^2 - (7)^2 = x^2 - 49$$

$$2. (a + 2b)(a - 2b) = (a)^2 - (2b)^2 = a^2 - 4b^2$$

$$3. (3x - 2y)(3x + 2y) = (3x)^2 - (2y)^2 = 9x^2 - 4y^2$$

In all the examples that we have looked at, whether they were special products or not, a single rule is sufficient.



When multiplying two multinomials together, we multiply each of the terms in the first multinomial times each of the terms in the second multinomial and then combine like terms.

### ■ Example 3-2 E

Perform the indicated multiplication and simplify.

$$\begin{aligned} 1. (3x - y)(2x + 3y) &= 6x^2 + 9xy - 2xy - 3y^2 \\ &= 6x^2 + 7xy - 3y^2 \end{aligned}$$

Distribute  
multiplication  
Combine like  
terms

$$\begin{aligned} 2. (a - 2)(2a^2 + 3a + 2) &= 2a^3 + 3a^2 + 2a - 4a^2 - 6a - 4 \\ &= 2a^3 - a^2 - 4a - 4 \end{aligned}$$

Distribute  
multiplication  
Combine like  
terms

**Note** Although there are three terms in the second parentheses, we still follow the procedure of every term in the first parentheses times every term in the second parentheses.

$$\begin{aligned} 3. (x - y)(x^2 + 3xy - y^2) &= x^3 + 3x^2y - xy^2 - x^2y - 3xy^2 + y^3 \\ &= x^3 + 2x^2y - 4xy^2 + y^3 \end{aligned}$$

Distribute  
multiplication

Combine like  
terms

4.  $(a + 6)(a - 2)(a - 1)$ . When there are three multinomials to be multiplied, we apply the associative property to multiply two of them together first and take that product times the third.

$$\begin{aligned} [(a + 6)(a - 2)](a - 1) &= [a^2 - 2a + 6a - 12](a - 1) \\ &= [a^2 + 4a - 12](a - 1) \\ &= a^3 - a^2 + 4a^2 - 4a - 12a + 12 \\ &= a^3 + 3a^2 - 16a + 12 \end{aligned}$$

### Mastery points

Can you

- Multiply a monomial and a multinomial?
- Multiply multinomials?
- Find the special products of the square of a binomial or the difference of two squares?

## Exercise 3-2

Perform the indicated multiplication and simplify. See examples 3-2 A, B, C, D, and E.

**Examples**  $3ab^2(2a - 3b)$

**Solutions**  $= 3ab^2 \cdot 2a - 3ab^2 \cdot 3b$   
 $= 6a^2b^2 - 9ab^3$

Distributive property  
 Multiply monomials

$(2x + y)(x - 3y)$

$= 2x \cdot x - 2x \cdot 3y + y \cdot x - y \cdot 3y$   
 $= 2x^2 - 6xy + xy - 3y^2$   
 $= 2x^2 - 5xy - 3y^2$

Distributive property  
 Multiply monomials  
 Combine like terms

1.  $2ab(a^2 - bc + c^2)$

4.  $-ab(a^4 - a^2b^2 - b^4)$

7.  $3ab(a^2 - 2ab - b^2)$

10.  $(x^2y)(x^2 + y^2)(xy^2)$

13.  $(y - 9)(y - 4)$

16.  $(b - 1)(b - 1)$

19.  $(a + 3)(a - 3)$

22.  $(3 - 2y)(2 - y)$

25.  $(3k + w)(k - 6w)$

28.  $(2a + 3b)^2$

31.  $(a + 4b)(a^2 - 2ab + b^2)$

34.  $(x - y)(x^2 - 2xy + y^2)$

37.  $(a - 6)(a - 2)(a + 1)$

40.  $(a + b)^3$

43.  $(a - 2b)^3$

2.  $6x(4y + 7z)$

5.  $-5ab^2(3a^2 - ab + 4b^2)$

8.  $(2x)(x - y + 5)(5y)$

11.  $(x + 3)(x + 4)$

14.  $(z + 7)(z - 11)$

17.  $(R - 3)^2$

20.  $(3x + 2)(x - 4)$

23.  $(7 + 2x)(2x - 7)$

26.  $(a - 6bc)(5a + 4bc)$

29.  $(2a + 3b)(2a - 3b)$

32.  $(x - 2y)(2x^2 - 3xy + y^2)$

35.  $(x^2 - 2x - 3)(x^2 + x + 4)$

38.  $(2b - 1)(b + 2)(2b + 1)$

41.  $(a - b)^3$

3.  $3a(5b^2 - 7c^2)$

6.  $6x^2(4x^2 - 2x + 3)$

9.  $(3a)(2a - b)(2b^2)$

12.  $(a + 5)(a - 3)$

15.  $(a + 1)(a + 1)$

18.  $(R + 2)(R - 2)$

21.  $(3a - 5)(2a - 7)$

24.  $(4r + 3)(r - 12)$

27.  $(a + 6b)^2$

30.  $(4x - y)(4x + y)$


33.  $(x + 4)(6x^2 - 3x + 7)$

36.  $(a^2 - 3a + 6)(a^2 + 2a - 5)$



39.  $(a - b)(a + b)(2a - 3b)$

42.  $(2a + b)^3$


44. The area of the shaded region between the two circles is  $\pi(R + r)(R - r)$ . Perform the indicated multiplication.



45. When squares of  $c$  units on a side are cut from the corners of a square sheet of metal  $x$  units on a side, and the metal sheet is then folded up into a tray, the volume is  $c(x - 2c)(x - 2c)$ . Perform the indicated multiplication.







# Extra Credit Rocks

Sign up for a Discover® Student Card today and enjoy:

- 0% Intro APR\* on Purchases for 6 Months
- No Annual Fee
- Easiest Online Account Management Options
- Full 5% *Cashback Bonus*®\* on Get More purchases in popular categories all year
- Up to 1% *Cashback Bonus*®\* on all your other purchases
- Unlimited cash rewards that never expire as long as you use your Card

**APPLY NOW**

**DISCOVER**  
CARD

\*View Discover® Card Rates, Fees, Rewards and Other Important Information.

**Review exercises**

Perform the indicated addition or subtraction. See sections 1-4 and 1-5.

1.  $(-3) + (-2)$

2.  $8 - (-4)$

3.  $(-7) - (-10)$

4.  $2 + (-5) + (-6)$

Simplify by using the properties of exponents. See section 3-1.

5.  $x^4 \cdot x^8$

6.  $(a^3)^5$

7.  $(3ab)^3$

8.  $(2x)^3$

**3-3 ■ Exponents—II****Fraction to a power property of exponents**

In section 3-1, we learned several useful properties of exponents. Now we shall learn several more.

Our next property of exponents can be derived from the definition of exponents. Consider the expression  $\left(\frac{a}{b}\right)^3$ .

$$\left(\frac{a}{b}\right)^3 = \frac{a}{b} \cdot \frac{a}{b} \cdot \frac{a}{b} = \frac{\overbrace{a \cdot a \cdot a}^{3 \text{ factors of } a}}{\underbrace{b \cdot b \cdot b}_{3 \text{ factors of } b}} = \frac{a^3}{b^3}$$

Thus

Fraction raised  
to a power

$$\left(\frac{a}{b}\right)^3 = \frac{a^3}{b^3}$$

Numerator raised to the power  
 Denominator raised to the power

**Fraction to a power property of exponents**

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}, b \neq 0$$

**Concept**

Whenever a fraction is raised to a power, the numerator and the denominator are *both* raised to that power.

**■ Example 3-3 A**

Perform the indicated operations and simplify.

1.  $\left(\frac{3}{4}\right)^3 = \frac{3^3}{4^3} = \frac{27}{64}$

2.  $\left(\frac{a}{b}\right)^5 = \frac{a^5}{b^5}$

3.  $\left(\frac{2a}{b}\right)^3 = \frac{(2a)^3}{b^3} = \frac{2^3 a^3}{b^3} = \frac{8a^3}{b^3}$



**Division of expressions with like bases**

Consider the expression

$$\frac{x^6}{x^2}$$

We can use the definition of exponents to write the fraction as

$$\frac{x^6}{x^2} = \frac{x \cdot x \cdot x \cdot x \cdot x \cdot x}{x \cdot x}$$

We reduce the fraction as follows:

$$\frac{x \cdot x \cdot x \cdot x \cdot x \cdot x}{x \cdot x} = \frac{x \cdot x \cdot x \cdot x}{1} = \frac{x^4}{1} = x^4$$

In our example, we reduced by two factors of  $x$ , leaving  $6 - 2 = 4$  factors of  $x$  in the numerator. Therefore

$$\frac{x^6}{x^2} = x^{6-2} = x^4$$

Thus we have the following property of exponents.

**Quotient property of exponents**

$$a^m \div a^n = \frac{a^m}{a^n} = a^{m-n}, a \neq 0$$

**Concept**

To divide quantities having *like* bases, subtract the exponent of the denominator from the exponent of the numerator to get the power of the given base in the quotient.

**Note** If the base  $a$  is zero,  $a = 0$ , we have an expression that has no meaning. Therefore  $a \neq 0$  indicates that we want our variables to assume no values that would cause the denominator to be zero.

**Example 3-3 B**

Simplify. Assume that no variable is equal to zero.

$$1. \frac{x^7}{x^5} = x^{7-5} = x^2$$

$$2. a^{11} \div a^4 = a^{11-4} = a^7$$

$$3. \frac{5^4}{5} = 5^{4-1} = 5^3 = 125$$

**Note** Remember that when we are dividing like bases, their exponents are subtracted, but *the base is not changed*.

$$4. \frac{a^5 \cdot a^2}{a^4} = \frac{a^{5+2}}{a^4} = \frac{a^7}{a^4} = a^{7-4} = a^3$$

$$5. \frac{x^3}{y^2} = \frac{x^3}{y^2}$$

**Note** In example 5, we cannot simplify. The bases must be the same in order to subtract exponents when we divide.

$$6. \frac{2^5 x^9 y^{15}}{2^3 x^5 y^{12}} = 2^{5-3} x^{9-5} y^{15-12} = 2^2 x^4 y^3 = 4x^4 y^3$$

► **Quick check** Simplify. Assume that no variable is equal to zero.  $a^{11} \div a^7$  ■

### Negative exponents

To this point, we have considered only those problems where the exponent of the numerator is greater than the exponent of the denominator. Consider the example

$$\frac{x^2}{x^6}$$

By the definition of exponents, this becomes

$$\frac{x^2}{x^6} = \frac{x \cdot x}{x \cdot x \cdot x \cdot x \cdot x \cdot x}$$

and reducing the fraction,

$$\frac{x \cdot x}{x \cdot x \cdot x \cdot x \cdot x \cdot x} = \frac{1}{x \cdot x \cdot x \cdot x} = \frac{1}{x^4}$$

Again, we reduced by two factors of  $x$ , leaving  $6 - 2 = 4$  factors of  $x$  in the denominator. Hence

$$\frac{x^2}{x^6} = \frac{1}{x^4}$$

However using the quotient property of exponents to carry out the division, we would have

$$\frac{x^2}{x^6} = x^{2-6} = x^{-4}$$

Since we should arrive at the same answer regardless of which procedure we use, then  $x^{-4}$  must be  $\frac{1}{x^4}$ , thus  $x^{-4} = \frac{1}{x^4}$ . This leads us to the definition of negative exponents.

### Definition of negative exponents

$$a^{-n} = \frac{1}{a^n}, a \neq 0$$

#### Concept

A negative exponent on any base (except zero) can be written as 1 over that base with a positive exponent.

### ■ Example 3-3 C

Write the following problems with positive exponents. Assume that no variable is equal to zero.

$$1. x^{-3} = \frac{1}{x^3}$$

Rewritten as 1 over  $x$  to the positive 3rd

$$2. a^{-9} = \frac{1}{a^9}$$

Rewritten as 1 over  $a$  to the positive 9th



**Note** From the definition of negative exponents, if a **factor** is moved from either the numerator to the denominator or from the denominator to the numerator, the sign of its exponent will change. The sign of the base will not be affected by this change.

**2. Alternative procedure**

$$\begin{aligned} a^{-9} &= \frac{a^{-9}}{1} \\ &= \frac{1}{a^9} \end{aligned}$$

Rewrite as a fraction

Sign of the exponent is changed as the **factor** is moved from the numerator to the denominator

$$\begin{aligned} 3. \frac{1}{b^{-4}} &= \frac{b^4}{1} \\ &= b^4 \end{aligned}$$

Sign of the exponent is changed as the **factor** is moved from the denominator to the numerator  
Standard form is to leave only positive exponents

$$\begin{aligned} 4. (-3)^{-3} &= \frac{1}{(-3)^3} \\ &= \frac{1}{-27} \text{ or } -\frac{1}{27} \end{aligned}$$

Sign of the exponent is changed as the factor is moved from the numerator to the denominator

Standard form

► **Quick check** Write  $b^{-2}$  with positive exponents. ■

### Zero as an exponent

Now consider the situation involving the division of like bases that are raised to the same power.

$$\frac{x^3}{x^3}, x \neq 0$$

By the definition of exponents, we have

$$\frac{x^3}{x^3} = \frac{x \cdot x \cdot x}{x \cdot x \cdot x} = \frac{1}{1} = 1$$

By the quotient property of exponents,

$$\frac{x^3}{x^3} = x^{3-3} = x^0$$

Since  $\frac{x^3}{x^3} = 1$  and  $\frac{x^3}{x^3} = x^0$ , then  $x^0$  must be equal to 1. This leads us to the definition of zero as an exponent.

#### Definition of zero as an exponent

$$a^0 = 1, a \neq 0$$

#### Concept

Any number other than zero raised to the zero power is equal to 1.

### ■ Example 3-3 D

Simplify. Assume that no variable is equal to zero.

1.  $b^0 = 1$
2.  $r^0 = 1$
3.  $5^0 = 1$
4.  $(-2)^0 = 1$
5.  $(a + b)^0 = 1$
6.  $3x^0 = 3 \cdot 1 = 3$
7.  $(3x)^0 = 1$

**Note** The exponent acts only on the symbol immediately to its left. In example 6, only the  $x$  is raised to the zero power. The exponent of 3 is understood to be 1. In example 7, the parentheses indicate that both the 3 and the  $x$  are raised to the zero power.

► **Quick check** Simplify. Assume that no variable is equal to zero.  $C^0$

### ■ Example 3-3 E

Simplify. Leave the answer with only positive exponents. Assume that no variable is equal to zero.

$$\begin{aligned} 1. \frac{x^5}{x^{11}} &= x^{5-11} \\ &= x^{-6} \\ &= \frac{1}{x^6} \end{aligned}$$

Division of like bases

Subtract exponents

Standard form

$$\begin{aligned} 2. a^{-7} \cdot a^5 &= a^{-7+5} \\ &= a^{-2} \\ &= \frac{1}{a^2} \end{aligned}$$

Multiplication of like bases

Add exponents

Standard form

$$\begin{aligned} 3. (b^{-2})^{-4} &= b^{(-2) \cdot (-4)} \\ &= b^8 \end{aligned}$$

Power of a power

Multiply exponents

$$\begin{aligned} 4. \frac{a^3b^5}{a^7b^2} &= a^{3-7}b^{5-2} \\ &= a^{-4}b^3 \\ &= \frac{b^3}{a^4} \end{aligned}$$

Division of like bases

Subtract exponents

Standard form

$$\begin{aligned} 5. \frac{a^3b^2c^4}{ab^5c^4} &= a^{3-1}b^{2-5}c^{4-4} \\ &= a^2b^{-3}c^0 \\ &= \frac{a^2 \cdot 1}{b^3} \\ &= \frac{a^2}{b^3} \end{aligned}$$

Division of like bases

Subtract the exponents

The  $a$ 's remain in the numerator, the  $b$ 's drop to the denominator, and  $c^0$  is 1

Standard form

$$\begin{aligned} 6. \frac{a^{-2}b^4}{a^{-5}b^6} &= a^{-2-(-5)}b^{4-6} \\ &= a^3b^{-2} \\ &= \frac{a^3}{b^2} \end{aligned}$$

Division of like bases

Subtract exponents

Standard form

► **Quick check** Simplify. Leave the answer with only positive exponents. Assume that no variable is equal to zero.  $\frac{b^4}{b^{10}}$



**Mastery points****Can you**

- Raise a fraction to a power?
- Perform division on expressions having like bases?
- Perform operations involving negative exponents?
- Perform operations involving zero as an exponent?

**Exercise 3-3**

Write each expression with only positive exponents. Assume that no variable is equal to zero. See examples 3-3 C and D.

**Examples**  $C^0$ **Solutions**  $= 1$  By definition is equal to 1 $b^{-2}$  $= \frac{1}{b^2}$  Rewritten as 1 over  $b$  to the positive 2nd

- |                 |                        |                         |                         |                  |
|-----------------|------------------------|-------------------------|-------------------------|------------------|
| 1. $x^0$        | 2. $(2y)^0$            | 3. $5a^0$               | 4. $7x^0$               | 5. $(3B)^0$      |
| 6. $S^{-2}$     | 7. $R^{-5}$            | 8. $(2x)^{-3}$          | 9. $(3P)^{-2}$          | 10. $4z^{-2}$    |
| 11. $9C^{-4}$   | 12. $\frac{5}{x^{-4}}$ | 13. $\frac{1}{2y^{-3}}$ | 14. $\frac{1}{3x^{-2}}$ | 15. $2x^{-4}y^2$ |
| 16. $x^{-2}y^4$ | 17. $p^0r^{-2}t^5$     | 18. $x^{-3}y^2z^{-4}$   |                         |                  |

Perform all indicated operations and leave your answer with only positive exponents. Assume that no variable is equal to zero. See examples 3-3 A, B, and E.

**Examples**  $a^{11} \div a^7$ **Solutions**  $= a^{11-7}$   
 $= a^4$  Division of like bases  
Subtract exponents $\frac{b^4}{b^{10}}$  $= b^{4-10}$   
 $= b^{-6}$   
 $= \frac{1}{b^6}$  Division of like bases  
Subtract exponents  
Standard form

- |                                    |                                   |                                   |                                  |                                   |
|------------------------------------|-----------------------------------|-----------------------------------|----------------------------------|-----------------------------------|
| 19. $\left(\frac{a}{b}\right)^6$   | 20. $\left(\frac{x}{y}\right)^4$  | 21. $\left(\frac{2}{3}\right)^3$  | 22. $\left(\frac{1}{2}\right)^4$ | 23. $\left(\frac{2x}{y}\right)^4$ |
| 24. $\left(\frac{2ab}{c}\right)^3$ | 25. $\left(\frac{3a}{b}\right)^3$ | 26. $x^{12} \div x^6$             | 27. $y^4 \div y^2$               | 28. $\frac{a^5}{a^3}$             |
| 29. $\frac{b^9}{b^7}$              | 30. $\frac{c^6}{c^9}$             | 31. $\frac{R^4}{R^8}$             | 32. $\frac{3^4}{3^2}$            | 33. $\frac{2^5}{2^3}$             |
| 34. $\frac{4^2}{4^5}$              | 35. $\frac{6}{6^3}$               | 36. $\frac{x^4x^3}{x^2}$          | 37. $\frac{y^5y}{y^2}$           | 38. $\frac{a^4a^2}{a^5}$          |
| 39. $\frac{a^4}{a^2a}$             | 40. $\frac{x^7}{x^2x^3}$          | 41. $\frac{y^3}{y^4y^5}$          | 42. $\frac{b^2}{bb^4}$           | 43. $\frac{a^7b^5}{a^4b^2}$       |
| 44. $\frac{x^9y^7}{x^4y}$          | 45. $\frac{2^3x^3y^7}{2xy^5}$     | 46. $\frac{3^3a^4b^5}{3^2a^2b^3}$ | 47. $\frac{3a^2b^5}{3^4a^5b^5}$  | 48. $\frac{5^2a^3b}{5^3a^7b^3}$   |

49.  $x^{-4}x^7$

54.  $x^5x^0x^{-2}$

59.  $\frac{3^{-2}}{3^{-5}}$

64.  $(y^{-3})^4$

69.  $(x^{-2})^0$

50.  $y^{-2}y^{10}$

55.  $a^0a^{-5}a^3$

60.  $\frac{2^{-6}}{2^{-3}}$

65.  $(a^{-2})^{-3}$

70.  $(b^0)^{-4}$

51.  $a^5a^{-11}$

56.  $a^{-7}a^4a^0$

61.  $(a^{-2})^3$

66.  $(z^{-4})^{-4}$

71.  $\frac{R^2S^{-4}}{R^{-3}S^5}$

52.  $R^{-2}R^{-5}$

57.  $(-5)^{-3}$

62.  $(b^4)^{-4}$

67.  $(x^0)^{-2}$

72.  $\frac{4y^{-3}}{4^{-1}y^2}$

53.  $x^{-2}x^4x^0$

58.  $(-2)^{-4}$

63.  $(x^5)^{-2}$

68.  $(a^{-3})^0$

73.  $\frac{4^{-1}a^{-2}b^3c^0}{2a^{-3}b^{-1}c^{-2}}$

**Review exercises**

Perform the indicated operations. See sections 1-4 to 1-8.

1.  $(-4) + (-6)$

2.  $(-3)(-7)$

3.  $(-4) - (-8)$

4.  $-4^2$

Simplify by using the properties of exponents. See section 3-1.

5.  $a^3a^5$

6.  $(x^3)^4$

7.  $xx^2x^3$

8.  $(2ab)^2$

**3-4 ■ Exponents—III****Properties and definitions of exponents**

The following is a summary of the properties and definitions of exponents that we have studied so far.

**Definitions**

$$a^n = \overbrace{a \cdot a \cdot a \cdots a}^{n \text{ factors}}, \text{ where } n \text{ is a positive integer}$$

$$a^{-n} = \frac{1}{a^n}, a \neq 0$$

$$a^0 = 1, a \neq 0$$

**Properties**

$$a^m \cdot a^n = a^{m+n}$$

$$(ab)^n = a^n b^n$$

$$(a^m)^n = a^{m \cdot n}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}, b \neq 0$$

$$a^m \div a^n = \frac{a^m}{a^n} = a^{m-n}, a \neq 0$$

The following examples illustrate some more problems in which more than one property of exponents is applied within the same problem.



### ■ Example 3-4 A

Simplify. Leave the answer with only positive exponents. Assume that no variable is equal to zero.

$$1. (2a^2b^3)^3 = 2^3(a^2)^3(b^3)^3$$

$$= 2^3a^6b^9$$

$$= 8a^6b^9$$

Each factor in the group is raised to the 3rd power

Power of a power, multiply exponents

8 is the standard form of  $2^3$

$$2. (5a^4b^2)^4 = 5^4(a^4)^4(b^2)^4$$

$$= 5^4a^{16}b^8$$

$$= 625a^{16}b^8$$

Each factor is raised to the power

Power of a power

Standard form

$$3. (3a^{-2}b^3)^{-3} = 3^{-3}(a^{-2})^{-3}(b^3)^{-3}$$

$$= 3^{-3}a^6b^{-9}$$

$$= \frac{a^6}{3^3b^9}$$

$$= \frac{a^6}{27b^9}$$

Each factor is raised to the power

Power of a power

The 3s and the b's drop to the denominator

Standard form

$$4. (-3a^2)(2ab^3)(-4a^3b^5)$$

$$= [(-3)(2)(-4)](a^2aa^3)(b^3b^5)$$

$$= 24a^6b^8$$

Multiply like bases using the commutative and associative properties

Signed numbers and multiplication of like bases, add exponents

$$5. (3a^4b)^2(3^2ab^5)^2 = 3^2(a^4)^2b^2 \cdot (3^2)^2a^2(b^5)^2$$

$$= 3^2a^8b^2 \cdot 3^4a^2b^{10}$$

$$= (3^2 \cdot 3^4)(a^8a^2)(b^2b^{10})$$

$$= 3^6a^{10}b^{12}$$

$$= 729a^{10}b^{12}$$

Group of factors to a power

Power of a power

Multiply like bases

Add exponents

Standard form

$$6. \left(\frac{2a^2b^3}{c^5}\right)^3 = \frac{(2a^2b^3)^3}{(c^5)^3}$$

$$= \frac{2^3(a^2)^3(b^3)^3}{(c^5)^3}$$

$$= \frac{8a^6b^9}{c^{15}}$$

Both the numerator and the denominator are raised to the 3rd power

Each factor in the numerator is raised to the 3rd power

Power of a power, multiply exponents

$$7. \frac{a^{-2}b^3}{a^{-4}b^6} = a^{(-2) - (-4)}b^{3 - 6}$$

$$= a^2b^{-3}$$

$$= \frac{a^2}{b^3}$$

Division of like bases

Subtract exponents

Standard form

$$8. \left(\frac{a^{-2}b}{c^3}\right)^{-2} = \frac{(a^{-2}b)^{-2}}{(c^3)^{-2}}$$

$$= \frac{(a^{-2})^{-2}b^{-2}}{(c^3)^{-2}}$$

$$= \frac{a^{(-2)(-2)}b^{-2}}{c^{(3)(-2)}}$$

$$= \frac{a^4b^{-2}}{c^{-6}}$$

$$= \frac{a^4c^6}{b^2}$$

Numerator and denominator are raised to the power

Numerator has a group of factors to a power

Power of a power

Multiply exponents

Standard form, factors raised to a negative power are moved to the other side of the fraction bar

► **Quick check** Simplify. Leave the answer with only positive exponents. Assume that no variable is equal to zero.  $(2a^{-2}b^3)^3$  ■

### Mastery points

Can you

- Apply the definitions and properties of exponents?

## Exercise 3-4

Simplify by using the properties and definitions of exponents. Leave the answer with only positive exponents. Assume that no variable is equal to zero. See example 3-4 A.

**Example**  $(2a^{-2}b^3)^3$

$$\begin{aligned} \text{Solution } &= 2^3(a^{-2})^3(b^3)^3 && \text{Groups of factors to a power} \\ &= 2^3a^{-6}b^9 && \text{Power of a power} \\ &= \frac{8b^9}{a^6} && \text{Standard form} \end{aligned}$$

- |   |   |  |  |
|---|---|--|--|
| 1. $(2a^2)^3$                               | 2. $(3x^4)^2$                                       | 3. $(2x^2y)^3$                                 | 4. $(4ab^3)^2$                                   |
| 5. $(x^4y^3z)^4$                            | 6. $(2a^5b^2c)^3$                                   | 7. $(5a^5b^2c^4)^2$                            | 8. $(a^3b^2)^3$                                  |
| 9. $(2a^2)^{-2}$                            | 10. $(5x^{-3})^{-2}$                                | 11. $(4^{-1}x^2)^{-2}$                         | 12. $(2a^2b^{-3})^{-2}$                          |
| 13. $(3xy^{-4})^{-3}$                       | 14. $(3x^{-2}y^{-3})^2$                             | 15. $(x^2y^{-5}z^3)^{-2}$                      | 16. $(a^{-3}b^2c^{-4})^{-3}$                     |
| 17. $(3x^2)(2x^0y^2)(x^5y)$                 | 18. $(a^2b)(-3a^0b^2)(a^3b)$                        | 19. $(-2x^2y)(3x^3y^2)(x^5y)$                  |  |
| 20. $(a^2bc)(-2a^2b^2c^2)(3abc)$            | 21. $(x^3yz^4)(-3xyz^2)(-2x^2yz)$                   | 22. $(a^2b)(-3b^2c^2)(2a^2c^2)$                |  |
| 23. $\left(\frac{2x}{y^2}\right)^3$         | 24. $\left(\frac{x^2y}{z^2}\right)^3$               | 25. $\left(\frac{3a^2c^0}{b^3}\right)^2$       | 26. $\left(\frac{x^3}{y^0z^4}\right)^3$          |
| 27. $\left(\frac{2x^2}{y^3}\right)^3$       | 28. $\left(\frac{ab^2}{c^4}\right)^4$               | 29. $\left(\frac{2x^3y^3}{z^5}\right)^2$       | 30. $\left(\frac{a^5bc^4}{d^2e}\right)^5$        |
| 31. $\frac{a^{-2}b^3}{a^3b^{-5}}$           | 32. $\frac{x^{-5}y^2}{x^3y^{-4}}$                   | 33. $\frac{3R^{-1}S^{-2}}{9R^{-3}S^2}$         | 34. $\frac{R^2S^{-4}}{R^{-3}S^5}$                |
| 35. $\frac{8a^{-2}b^{-5}}{2a^{-1}b^4}$      | 36. $\frac{2x^{-1}y^{-2}}{3x^{-2}y^2}$              | 37. $\frac{6R^{-2}S^0}{2R^2S^{-3}}$            | 38. $\frac{2a^{-1}b^0c^2}{5a^3b^{-1}c^{-3}}$     |
| 39. $(a^2b^3)^3(ab^2)^4$                    | 40. $(xy^2)^3(x^2y^2)^2$                            | 41. $(2a^3)^2(2a^2)^3$                         | 42. $(3x^5)^3(3x^3)^2$                           |
| 43. $(2x^2y)^3(2x^4y^5)^2$                  | 44. $(3r^2s^4)^3(r^5s^6)^2$                         | 45. $\left(\frac{xy^{-2}}{z^{-4}}\right)^{-1}$ | 46. $\left(\frac{x^{-3}y}{z^5}\right)^{-2}$      |
| 47. $\left(\frac{2a^{-3}}{b^5}\right)^{-2}$ | 48. $\left(\frac{4^{-1}a^{-2}}{b^{-5}}\right)^{-2}$ | 49. $\left(\frac{ab^{-2}}{c^{-1}}\right)^{-3}$ | 50. $\left(\frac{2^{-2}x^3}{y^{-2}}\right)^{-3}$ |

## Review exercises

Perform the indicated multiplication. See section 1-2.

- |                        |                        |                        |
|------------------------|------------------------|------------------------|
| 1. $(6.2) \cdot (5.7)$ | 2. $(2.8) \cdot (3.7)$ | 3. $(1.9) \cdot (8.8)$ |
| 4. $(4.2) \cdot (6.9)$ | 5. $(9.9) \cdot (1.9)$ | 6. $(7.5) \cdot (6.6)$ |



## 3-5 ■ Scientific notation

**Scientific notation**

An important use of integer exponents is in scientific, engineering, and technical fields where we deal with very large or very small numbers. For example, the mass of a hydrogen atom is 0.000 000 000 000 000 000 000 001 67 gram; the mass of an electron is 0.000 000 000 000 000 000 000 000 000 91 gram; the half-life of lead-204 is 14,000,000,000,000,000,000 years. To work with such numbers on the calculator, they must often be entered in scientific notation. We define the scientific notation of a positive number  $X$  to be the product

$$X = a \times 10^n$$

where  $1 \leq a < 10$  and  $n$  is an integer. To achieve this form of the decimal number  $X$ , use the following steps.

**Scientific notation**

- Step 1** Move the decimal point to a position immediately following the first nonzero digit in  $X$ .
- Step 2** Count the number of places the decimal point has been moved. This is the power,  $n$ , to which 10 is raised.
- Step 3** If
- the decimal point is moved to the *left*,  $n$  is *positive*.
  - the decimal point is moved to the *right*,  $n$  is *negative*.
  - the decimal point already follows the first nonzero digit,  $n$  is zero.

■ **Example 3-5 A**

Express the following numbers in scientific notation.

1. 250

$$250 = \underbrace{2.50}_{\text{decimal point moved 2 places right}} \times 10^2 = 2.5 \times 10^2$$

2. 45,000,000

$$45,000,000 = \underbrace{4.5000000}_{\text{decimal point moved 7 places right}} \times 10^7 = 4.5 \times 10^7$$

3. 5

$$5 = 5 \times 10^0$$

4. 0.000152

$$0.000152 = \underbrace{0.000152}_{\text{decimal point moved 4 places right}} \times 10^{-4} = 1.52 \times 10^{-4}$$

**Note** To write a negative number in scientific notation, we use the same procedure as for a positive number except that a negative sign,  $-$ , is placed in front of  $a$ .

**Example**

To write  $-0.0234$  in scientific notation, we proceed as follows:

$$-0.0234 = \underbrace{-0.0234}_{\text{decimal point moved 2 places right}} \times 10^{-2} = -2.34 \times 10^{-2}$$

► **Quick check** Express the following numbers in scientific notation.

4,380       $-0.00592$

**Standard form**

Sometimes it is necessary to convert a number in scientific notation to its standard form. To do this, we apply the rules in reverse.

**Standard form**

When the power of 10 is

1. *positive*, the decimal point is moved to the *right*  $n$  places.
2. *negative*, the decimal point is moved to the *left*  $n$  places.
3. *zero*, the decimal point is not moved.

**■ Example 3-5 B**

Express the following numbers in standard form.

1.  $1.45 \times 10^4$

Since the exponent of 10 is positive 4, we move the decimal point 4 places to the *right* to get

$$1.45 \times 10^4 = \underbrace{1.4500}_{\text{move decimal 4 places right}} = 14,500$$

2.  $5.23 \times 10^{-3}$

The *negative* exponent,  $-3$ , tells us to move the decimal point 3 places to the *left* to get

$$5.23 \times 10^{-3} = \underbrace{0.00523}_{\text{move decimal 3 places left}} = 0.00523$$

**Note** In each example, it was necessary to insert zeros to properly locate the decimal point.

3.  $-4.07 \times 10^{-2}$

With a negative exponent,  $-2$ , move the decimal point 2 places to the *left* to get

$$-4.07 \times 10^{-2} = \underbrace{-0.0407}_{\text{move decimal 2 places left}} = -0.0407$$

**Note** The negative sign preceding the number is carried along into the standard form.

► **Quick check** Express the following numbers in standard form.

$$9.98 \times 10^{-4} \qquad -5.63 \times 10^4$$

**Computation using scientific notation**

Scientific notation can be used to simplify numerical calculations when the numbers are very large or very small. We first change the numbers to scientific notation and use the properties of exponents to help perform the indicated operations.

**■ Example 3-5 C**

Perform the indicated operations using scientific notation.

1.  $(349,000,000)(0.0816)$

$$= (3.49 \times 10^8)(8.16 \times 10^{-2})$$

$$= (3.49 \cdot 8.16) \times (10^8 \cdot 10^{-2})$$

$$= 28.4784 \times 10^6$$

$$= 28,478,400$$

Scientific notation

Commutative and associative properties

Multiply

Standard form



$$\begin{aligned}
 2. \quad & \frac{(102,000,000)(0.00105)}{(1,190)(0.012)} \\
 &= \frac{(1.02 \times 10^8)(1.05 \times 10^{-3})}{(1.19 \times 10^3)(1.2 \times 10^{-2})} && \text{Scientific notation} \\
 &= \frac{(1.02)(1.05)10^8 \cdot 10^{-3}}{(1.19)(1.2)10^3 \cdot 10^{-2}} && \text{Commutative and associative properties} \\
 &= \frac{(1.02)(1.05)}{(1.19)(1.2)} \times 10^4 && \text{Properties of exponents} \\
 &= 0.75 \times 10^4 && \text{Multiplication and division} \\
 &= 7,500 && \text{Standard form}
 \end{aligned}$$

**Mastery points***Can you*

- Express a number in scientific notation?
- Convert a number from scientific notation to standard form?
- Do computations using scientific notation?

**Exercise 3–5**

Express the following numbers in scientific notation. See example 3–5 A.

**Examples** 4,380**Solutions**  $= 4.38 \times 10^3$ Three places to the left,  
exponent is 3

–0.00592

 $= -5.92 \times 10^{-3}$ Three places to the right,  
exponent is –3

- |                |                    |               |                 |                    |
|----------------|--------------------|---------------|-----------------|--------------------|
| 1. 255         | 2. 65,000,000      | 3. 12,345     | 4. 14,800       | 5. 155,000         |
| 6. 14.36       | 7. 855.076         | 8. 1,570.7    | 9. 1,007,600    | 10. 6,000,736      |
| 11. 0.00012    | 12. 0.0863         | 13. 0.0000081 | 14. 0.0000147   | 15. 0.0007         |
| 16. 0.12079    | 17. 0.000000000094 | 18. –456      | 19. –4,500      | 20. –0.00087       |
| 21. –5,850,000 | 22. –0.0567        | 23. –45.78    | 24. –34,000,000 | 25. –0.00000002985 |

Convert the following numbers in scientific notation to their standard form. See example 3–5 B.

**Examples**  $9.98 \times 10^{-4}$ **Solutions**  $= 0.000998$ Exponent is –4, move 4 places  
to the left $-5.63 \times 10^4$  $= -56,300$ Exponent is 4, move 4 places  
to the right

- |                          |                            |                            |                            |
|--------------------------|----------------------------|----------------------------|----------------------------|
| 26. $2.07 \times 10^3$   | 27. $4.99 \times 10^7$     | 28. $5.061 \times 10^5$    | 29. $7.23 \times 10^0$     |
| 30. $1.073 \times 10^4$  | 31. $4.2 \times 10^{-3}$   | 32. $7.611 \times 10^{-7}$ | 33. $1.47 \times 10^{-6}$  |
| 34. $5.0 \times 10^{-2}$ | 35. $7.89 \times 10^{-4}$  | 36. $-2.3 \times 10^5$     | 37. $-4.82 \times 10^{-9}$ |
| 38. $-2.61 \times 10^2$  | 39. $-4.92 \times 10^{-6}$ | 40. $-9.3 \times 10^8$     |                            |

Express the following numbers in scientific notation or in standard form. See examples 3–5 A and B.

41. A millicron equals 0.000000001 of a meter. Write this number in scientific notation.
42. The speed of light is approximately 30,000,000,000 centimeters per second. Write this in scientific notation.



## Student Loans for up to **\$40,000** per year\*

Defer payments until after graduation.\*\*  
Fast preliminary approval, usually in minutes.

 **Apply Now!**  
Go to [gmacbankfunding.com](http://gmacbankfunding.com)

## Apply online in as little as 15 minutes

- Loans up to \$40,000 per academic year\*
- Good for tuition and other educational expenses: books, fees, a laptop, room and board, travel home, etc.
- Get a check in as few as 5 business days
- Start payments now or up to six months after graduation\*\*
- Reduce your interest rate by as much as 0.50% with automatic payments\*\*\*

All loans are subject to application and credit approval.

\* Undergraduate and graduate borrowers may borrow annually up to the lesser of the cost of attendance or \$30,000 (\$40,000 for certain schools where it has been determined that the annual cost of attendance exceeds \$30,000). Borrowers in the Continuing Education loan program may borrow annually up to \$30,000.

\*\* Undergraduate students may choose to defer repayment until six months after graduation or ceasing to be enrolled at least half time in school. Interest only and immediate repayment options also available.

\*\*\* A 0.25% interest rate reduction is available for borrowers who elect to have monthly principal and interest payments transferred electronically from a savings or checking account. The interest rate reduction will begin when automatic principal and interest payments start, and will remain in effect as long as automatic payments continue without interruption. This reduced interest rate will return to contract rate if automatic payments are cancelled, rejected or returned for any reason. Upon request, borrowers are also entitled to an additional 0.25% interest rate reduction if (1) the first 36 payments of principal and interest are paid on time, and (2) at any time prior to the 36th on time payment, the borrower who receives the monthly bill elects to have monthly principal and interest payments transferred electronically from a savings or checking account, and continues to make such automatic payments through the 36th payment. This reduced interest rate will not be returned to contract rate if, after receiving the benefit, the borrower discontinues automatic electronic payment. The lender and servicer reserve the right to modify or discontinue borrower benefit programs (other than the co-signer release benefit) at any time without notice.

GMAC Bank Member FDIC

### Chapter 3 summary

1. In the expression  $x^6$ ,  $x$  is called the **base** and 6 the **exponent**.

$n$  factors

2.  $a^n = \overbrace{a \cdot a \cdot a \cdots a}^n$ , where  $n$  is a positive integer.

3. **Properties and definitions of exponents**

$$a^m \cdot a^n = a^{m+n}$$

$$(ab)^n = a^n b^n$$

$$(a^m)^n = a^{m \cdot n}$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}, b \neq 0$$

$$a^m \div a^n = \frac{a^m}{a^n} = a^{m-n}, a \neq 0$$

$$a^{-n} = \frac{1}{a^n}, a \neq 0$$

$$a^0 = 1, a \neq 0$$

4. When *multiplying* two multinomials, we multiply each term in the first multinomial by each term in the second multinomial. We then combine like terms.

5. Three **special products** are:

$$(a + b)^2 = a^2 + 2ab + b^2$$

$$(a - b)^2 = a^2 - 2ab + b^2$$

$$(a + b)(a - b) = a^2 - b^2$$

6. The scientific notation of a positive number  $X$  is  $X = a \times 10^n$ , where  $1 \leq a < 10$  and  $n$  is an integer.

### Chapter 3 error analysis

1. Exponents

Example:  $xy^3 = x^3y^3$

Correct answer:  $xy^3 = xy^3$

What error was made? (see page 127)

2. Multiplication of like bases

Example:  $4^2 \cdot 4^3 = 16^5$

Correct answer:  $4^2 \cdot 4^3 = 4^5$

What error was made? (see page 129)

3. Power of a power

Example:  $(a^3)^2 = a^5$

Correct answer:  $a^6$

What error was made? (see page 130)

4. Multiplying unlike bases

Example:  $x^3 \cdot y = (xy)^4$

Correct answer:  $x^3 \cdot y = x^3y$

What error was made? (see page 131)

5. Product of a monomial and a multinomial

Example:  $-x(x^2 - 2x + 1) = x^3 - 2x^2 + x$

Correct answer:  $-x^3 + 2x^2 - x$

What error was made? (see page 134)

6. Squaring a binomial

Example:  $(4x + y)^2 = (4x)^2 + (y)^2 = 16x^2 + y^2$

Correct answer:  $16x^2 + 8xy + y^2$

What error was made? (see page 136)

7. Dividing like bases

Example:  $\frac{x^3}{x} = x^4$

Correct answer:  $x^2$

What error was made? (see page 140)

8. Negative exponents

Example:  $\frac{1}{a^{-3}} = -a^3$

Correct answer:  $\frac{1}{a^{-3}} = a^3$

What error was made? (see page 141)

9. Zero exponent

Example:  $(x + y)^0 = x^0 + y^0 = 1 + 1 = 2$

Correct answer:  $(x + y)^0 = 1$

What error was made? (see page 143)

10. Evaluate absolute value

Example:  $-|-4| = 4$

Correct answer:  $-|-4| = -4$

What error was made? (see page 31)

### Chapter 3 critical thinking

Given the number  $52^2$ , determine a method by which you can square the 52 mentally.

**Chapter 3 review****[3-1]**

Simplify and leave the answers with only positive exponents.

- |                        |                            |                       |
|------------------------|----------------------------|-----------------------|
| 1. $a^5 \cdot a^7$     | 2. $a \cdot a^4 \cdot a^9$ | 3. $4^3 \cdot 4^2$    |
| 4. $(xy)^4$            | 5. $(a^3)^5$               | 6. $(5ab^3)(4a^3b^2)$ |
| 7. $(3x^2y^3)(2xy^4)$  | 8. $(-5x^2)(3x^3)$         | 9. $(2a^2b)(3ab^4)$   |
| 10. $(5x^2y)(2x^3y^4)$ | 11. $(-3a^2b^3)(2a^4b^7)$  |                       |

**[3-2]**

Perform the indicated multiplication and simplify.

- |                         |                                 |                               |
|-------------------------|---------------------------------|-------------------------------|
| 12. $5x(3x - 2y)$       | 13. $-3a^2b(2a^2 - 3ab + 4b^2)$ | 14. $(5x)(3x - y)(2y^2)$      |
| 15. $(x + 3)(x - 4)$    | 16. $(x + 5)^2$                 | 17. $(a - 7)(a + 7)$          |
| 18. $(5x - y)(3x + 2y)$ | 19. $(x - 2y)(x^2 + 3xy + y^2)$ | 20. $(3a - b)(a + b)(a - 2b)$ |
| 21. $(2a + b)^3$        |                                 |                               |

**[3-3]**

Simplify and leave the answers with only positive exponents.

- |                                    |                               |                                 |                                  |
|------------------------------------|-------------------------------|---------------------------------|----------------------------------|
| 22. $\frac{b^5}{b^7}$              | 23. $5a^{-2}$                 | 24. $a^{-5} \cdot a^9$          | 25. $\frac{x^3x^2}{x^8}$         |
| 26. $(3a^2b)^0$                    | 27. $5x^{-3}y^{-2}$           | 28. $\frac{a^{-4}}{a^{-7}}$     | 29. $\left(\frac{a}{b}\right)^5$ |
| 30. $\left(\frac{2yz}{x}\right)^2$ | 31. $\frac{2a^2b^4}{2^3a^5b}$ | 32. $\frac{a^5b^{-2}}{a^{-4}b}$ |                                  |

**[3-4]**

Simplify and leave the answers with only positive exponents.

- |   |  |  |   |
|---|--|--|---|
| 33. $(2a^2b^3)^3$                               | 34. $(3^3x^4y^5)^4$                      | 35. $(2xy^{-3})^{-2}$                  | 36. $\frac{2x^{-1}y^0z^3}{4x^{-2}y^{-3}}$ |
| 37. $\frac{8a^{-5}b^{-4}c^0}{4a^{-7}b^2c^{-3}}$ | 38. $(x^5y^4)^4(2x^2y^3)^3$              | 39. $(2a^2b)^3(3a^4)^2$                |   |
| 40. $\left(\frac{a^3b^0}{c^4}\right)^3$         | 41. $\left(\frac{3a^3b^2}{c^5}\right)^2$ | 42. $\left(\frac{2xy^4}{z^6}\right)^5$ |   |

**[3-5]**

Express the following numbers in scientific notation.

- |                     |             |                 |
|---------------------|-------------|-----------------|
| 43. 1,840           | 44. 0.00157 | 45. 107,000,000 |
| 46. 849,000,000,000 | 47. -37.5   | 48. -0.00543    |

Express the following numbers in standard form.

- |                            |                           |                         |
|----------------------------|---------------------------|-------------------------|
| 49. $5.04 \times 10^5$     | 50. $6.39 \times 10^{-3}$ | 51. $-5.96 \times 10^2$ |
| 52. $-8.86 \times 10^{-3}$ | 53. $7.35 \times 10^{-7}$ | 54. $8.12 \times 10^8$  |

Perform the indicated operations using scientific notation. Leave the answer in scientific notation.

- |                                      |                                    |
|--------------------------------------|------------------------------------|
| 55. $(456,000,000) \cdot (0.000587)$ | 56. $(0.0000183) \cdot (0.000846)$ |
| 57. $(756,000) \div (105,000,000)$   | 58. $(0.00525) \div (42,000)$      |



**Chapter 3 cumulative test**

Determine if the following statements are true or false.

[1-3] 1.  $|-10| < 0$

[1-8] 2.  $-3^2 = (-3)^2$

[1-3] 3.  $|-3| < |-7|$

Perform the indicated operations, if possible, and simplify.

[1-7] 4.  $\frac{(8)}{(-4)}$

[1-7] 5.  $\frac{(-9)}{0}$

[1-5] 6.  $(2) - (-6)$

[1-6] 7.  $(-2)(4)(0)(-4)$

[1-8] 8.  $(-2)^4$

[1-8] 9.  $48 - 24 \div 8 - 3 - 2^2$

[2-3] 10.  $(3x^2y - 4xy + 2xy^2) - (2x^2y - 4xy + 3x^2y^2)$

[1-6] 11.  $(5)(-2)(4)$

[1-7] 12.  $\frac{0}{-3}$

[3-2] 13.  $(3x - y)^2$

[1-6] 14.  $(2)(-7)(0)(3)$

[3-4] 15.  $(2a^2b^5)^2$

[1-6] 16.  $-5^2$

[1-8] 17.  $10 - 10 \div 10 \cdot 10 - 10 + 10$

[3-2] 18.  $(3x - 2y)(3x + 2y)$

[3-1] 19.  $(3x^2y)(2xy^4)$

[1-8] 20.  $2[5(7 - 4) - 6 + 4]$

[1-7] 21.  $\frac{(-6)(-4)}{(5)(0)}$

[2-3] 22.  $(3a - 2b) - [5a - (4b + 6a)]$

[3-2] 23.  $(x + 1)(x^2 - x - 1)$

[3-3] 24.  $x^{-3}x^5x^0$

[3-3] 25.  $\frac{a^{-5}}{a^{-9}}$

Find the solution set.

[2-6] 26.  $3x - 4 = x + 10$

[2-6] 27.  $2(x - 4) + 7 = 8x - 11$

[2-6] 28.  $\frac{2}{3}x + 4 = \frac{5}{6}$

Find the solution.

[2-9] 29.  $8 - 3x < 9$

[2-9] 30.  $6x + 5 - 4 > 2$

[2-6] 31.  $3 - 2x = 6$

[2-9] 32.  $-9 \leq 2x + 7 \leq 5$

[2-8] 33. If a number is decreased by 17 and that result is then divided by 5, the answer is 16. Find the number.

[2-8] 34. One-third of a number is 12 less than one-half of the number. Find the number.

[2-8] 35. Brenda invested part of \$30,000 at 8% and the rest at 7%. If her income for one year from the 8% investment was \$675 more than that from the 7% investment, how much was invested at each rate?


## Review exercises

1. -16 2. 16 3. -16 4. 16 5.  $x^5$  6. (let  $x$  = the number)  $x^3$  7. (let  $x$  = the number)  $x^2$  8.  $xy$

## Chapter 2 review

1. 3 2. 1 3. 2 4. 2 5. polynomial 6. polynomial  
7. polynomial 8. not a polynomial because a variable is in the denominator 9.  $5x$  10.  $y - 7$  11.  $z + 4$   
12. (let  $x$  = the number)  $2x + 6$  13. 1 14. -1 15. 72  
16. -4 17. 4 18. 7 19. a. 3 b.  $\frac{189}{4}$  20. 1,040  
21.  $4x^2 - 3x + 3$  22.  $-a^2 + a + 11$  23.  $-6a^2$   
24.  $5x^3 - 7xy^2 - 2y^3 - 4x^2$  25.  $-11ab + 7b^2c + 11bc$   
26.  $-6y + 1$  27.  $5ab + 3ac - 4bc$  28. -5 29.  $7x + y$   
30.  $2x + 5y$  31.  $-6x + 4y$  32.  $9a - 10b$  33. true  
34. false 35. false 36. true 37. {7} 38. {21}  
39. {-11} 40. {-6} 41. {4} 42. {19} 43. {-2}  
44. {9} 45. {3} 46. {3} 47. {-7} 48. {-7} 49. {12}  
50. {14} 51. {15} 52. {21} 53.  $\left\{\frac{21}{4}\right\}$  54. {-18}  
55.  $\left\{-\frac{9}{2}\right\}$  56. {35} 57. {0} 58. {4} 59. {6} 60. {-8}  
61. {3} 62.  $\left\{\frac{14}{3}\right\}$  63.  $\left\{\frac{7}{3}\right\}$  64. {3} 65.  $\left\{\frac{1}{2}\right\}$   
66.  $\left\{-\frac{22}{3}\right\}$  67.  $\left\{-\frac{1}{4}\right\}$  68.  $\left\{\frac{7}{13}\right\}$  69.  $\left\{-\frac{5}{2}\right\}$  70. {-14}  
71. {-24} 72.  $\left\{\frac{7}{4}\right\}$  73. {0} 74.  $\left\{\frac{1}{5}\right\}$  75.  $\left\{-\frac{7}{4}\right\}$   
76. {2} 77.  $a = \frac{F}{m}$  78.  $I = \frac{E}{R}$  79.  $P = \frac{k}{V}$   
80.  $g = V - k - t$  81.  $c = \frac{2A - bh}{h}$  82.  $x = \frac{4y}{3}$   
83. 64 and 41 84. 36 85. 45 86. \$11,000 at 8%;  
\$9,000 at 7% 87. \$12,000 at 12%; \$13,000 at 19%


88.  $x > 4$ ;




90.  $x > -7$ ;




92.  $x < 2$ ;




94.  $x > \frac{9}{2}$ ;




89.  $x \leq 3$ ;




91.  $x < -4$ ;




93.  $x > \frac{15}{7}$ ;




95.  $x \geq -\frac{6}{5}$ ;




96.  $x \leq \frac{2}{9}$ ;




97.  $-\frac{11}{5} < x < \frac{3}{5}$ ;




98.  $-1 < x \leq \frac{1}{5}$ ;



99.  $\frac{1}{2} < x < \frac{9}{4}$ ;



100.  $-\frac{13}{3} \leq x \leq -\frac{1}{3}$ ;



## Chapter 2 cumulative test

1. -12 2. 4 3. 3 4. undefined 5. -25 6.  $\frac{1}{2}$   
7.  $-\frac{1}{6}$  8. 0 9. 19.78 10. 41 11. 38 12. -24  
13. 26 14. 6 15. 20 16.  $4x$  17.  $2x^2y^2 + 3xy$   
18.  $a + 3b$  19.  $4x^2y - 7xy^2$  20.  $a^3 + 2a^2 + a - 1$   
21.  $-3x + 4y$  22.  $4a + 2b$  23. 64 24. -25 25. 234  
26. 2 27. 4 28. 42 29. 51 mph 30.  $x - y$   
31. (let  $x$  = the number)  $x + 6$  32.  $10d + 5n + c$  33.  $\left\{\frac{5}{3}\right\}$   
34. {0} 35.  $\left\{-\frac{19}{2}\right\}$  36.  $\left\{\frac{17}{16}\right\}$  37.  $\left\{\frac{30}{11}\right\}$  38.  $x \leq -6$   
39.  $x < -7$  40.  $x > \frac{8}{5}$  41.  $-2 < x < 4$   
42.  $-1 \leq x \leq 6$  43.  $b = P - a - c$  44.  $y = \frac{x - az}{a}$   
45. \$6,000 at 6%; \$4,000 at 5% 46. 14, 16, 18 47. 12  
48. \$6,500 at 12% profit; \$10,500 at 19% loss 49.  $x \leq 6$

## Chapter 3

## Exercise 3-1

## Answers to odd-numbered problems

1.  $a^5$  3.  $(-2)^4$  5.  $x^6$  7.  $(xy)^4$  9.  $(x - y)^3$  11.  $xxxx$   
13.  $(-2)(-2)(-2)$  15.  $5 \cdot 5 \cdot 5$  17.  $(4y)(4y)(4y)(4y)$   
19.  $(x - y)(x - y)$  21.  $x^{11}$  23.  $R^3$  25.  $a^9$  27.  $5^5 = 3,125$   
29.  $4^7 = 16,384$  31.  $(x - 2y)^{10}$  33.  $(a - b)^{11}$  35.  $x^4y^4$   
37.  $64x^3y^3z^3$  39.  $x^{15}$  41.  $b^{25}$  43.  $6x^4y^3$  45.  $a^7b^5$   
47.  $30x^5$  49.  $12a^7b^7c$  51.  $12a^5b^3$  53.  $-6a^3b^5$   
55. a.  $V = 5^3$ , 125 cubic units b.  $V = 4^3$ , 64 cubic units  
c.  $V = 6^3$ , 216 cubic units 57.  $s = \frac{1}{2}gt^2$  59.  $V = \frac{4}{3}\pi r^3$   
61.  $m^4 - 8$  63.  $2x^2 - y^3$  65.  $\frac{p^3}{q^2}$



## Solutions to trial exercise problems

23.  $R^2 \cdot R = R^2 \cdot R^1 = R^{2+1} = R^3$  27.  $5^2 \cdot 5^3 = 5^{2+3} = 5^5 = 3,125$  43.  $(2xy^2)(3x^3y) = 2 \cdot 3 \cdot xx^3y^2y = 6x^{1+3}y^{2+1} = 6x^4y^3$   
 55a.  $V = e^3$ , then  $V = (5)^3 = 5 \cdot 5 \cdot 5 = 25 \cdot 5 = 125$  cubic units  
 60. The cube of Johnny's age is given as  $n^3$ . Since his mother is 6 years more than the cube of Johnny's age, we add 6, giving  $n^3 + 6$ .

## Review exercises

1.  $9a$  2.  $8x$  3.  $6ab$  4.  $7xy$  5.  $7a^2 + 5a$  6.  $5x^2 + 5x$   
 7.  $x^2y + 7xy^2$  8.  $3ab^2 + 2a^2b$

## Exercise 3-2

## Answers to odd-numbered problems

1.  $2a^3b - 2ab^2c + 2abc^2$  3.  $15ab^2 - 21ac^2$  5.  $-15a^3b^2 + 5a^2b^3 - 20ab^4$  7.  $3a^3b - 6a^2b^2 - 3ab^3$  9.  $12a^2b^2 - 6ab^3$   
 11.  $x^2 + 7x + 12$  13.  $y^2 - 13y + 36$  15.  $a^2 + 2a + 1$   
 17.  $R^2 - 6R + 9$  19.  $a^2 - 9$  21.  $6a^2 - 31a + 35$   
 23.  $4x^2 - 49$  25.  $3k^2 - 17kw - 6w^2$  27.  $a^2 + 12ab + 36b^2$   
 29.  $4a^2 - 9b^2$  31.  $a^3 + 2a^2b - 7ab^2 + 4b^3$   
 33.  $6x^3 + 21x^2 - 5x + 28$  35.  $x^4 - x^3 - x^2 - 11x - 12$   
 37.  $a^3 - 7a^2 + 4a + 12$  39.  $2a^3 - 3a^2b - 2ab^2 + 3b^3$   
 41.  $a^3 - 3a^2b + 3ab^2 - b^3$  43.  $a^3 - 6a^2b + 12ab^2 - 8b^3$   
 45.  $cx^2 - 4c^2x + 4c^3$

## Solutions to trial exercise problems

8.  $(2x)(x - y + 5)(5y) = [(2x)(x - y + 5)](5y)$   
 $= [2x \cdot x - 2x \cdot y + 2x \cdot 5](5y) = [2x^2 - 2xy + 10x](5y)$   
 $= 2x^2 \cdot 5y - 2xy \cdot 5y + 10x \cdot 5y = 10x^2y - 10xy^2 + 50xy$   
 17.  $(R - 3)^2$  is a special product.  $(R - 3)^2$   
 $= (R)^2 + [2 \cdot R \cdot (-3)] + (-3)^2 = R^2 - 6R + 9$   
 18.  $(R + 2)(R - 2)$  is a special product.  $(R + 2)(R - 2)$   
 $= (R)^2 - (2)^2 = R^2 - 4$  31.  $(a + 4b)(a^2 - 2ab + b^2)$   
 $= a^3 - 2a^2b + ab^2 + 4a^2b - 8ab^2 + 4b^3 = a^3 + 2a^2b - 7ab^2 + 4b^3$  37.  $(a - 6)(a - 2)(a + 1) = [(a - 6)(a - 2)](a + 1)$   
 $= [a^2 - 2a - 6a + 12](a + 1) = [a^2 - 8a + 12](a + 1)$   
 $= a^3 + a^2 - 8a^2 - 8a + 12a + 12 = a^3 - 7a^2 + 4a + 12$   
 40.  $(a + b)^3 = (a + b)(a + b)(a + b) = [(a + b)(a + b)](a + b)$   
 $= [a^2 + ab + ab + b^2](a + b) = [a^2 + 2ab + b^2](a + b)$   
 $= a^3 + a^2b + 2a^2b + 2ab^2 + ab^2 + b^3 = a^3 + 3a^2b + 3ab^2 + b^3$

## Review exercises

1.  $-5$  2.  $12$  3.  $3$  4.  $-9$  5.  $x^{12}$  6.  $a^{15}$  7.  $27a^3b^3$   
 8.  $8x^3$

## Exercise 3-3

## Answers to odd-numbered problems

1.  $1$  3.  $5$  5.  $1$  7.  $\frac{1}{R^5}$  9.  $\frac{1}{9p^2}$  11.  $\frac{9}{C^4}$  13.  $\frac{y^3}{2}$   
 15.  $\frac{2y^2}{x^4}$  17.  $\frac{t^5}{r^2}$  19.  $\frac{a^6}{b^6}$  21.  $\frac{8}{27}$  23.  $\frac{16x^4}{y^4}$  25.  $\frac{27a^3}{b^3}$   
 27.  $y^2$  29.  $b^2$  31.  $\frac{1}{R^4}$  33.  $4$  35.  $\frac{1}{36}$  37.  $y^4$  39.  $a$   
 41.  $\frac{1}{y^6}$  43.  $a^3b^3$  45.  $4x^2y^2$  47.  $\frac{1}{27a^3}$  49.  $x^3$  51.  $\frac{1}{a^6}$   
 53.  $x^2$  55.  $\frac{1}{a^2}$  57.  $-\frac{1}{125}$  59.  $27$  61.  $\frac{1}{a^6}$  63.  $\frac{1}{x^{10}}$   
 65.  $a^6$  67.  $1$  69.  $1$  71.  $\frac{R^5}{S^9}$  73.  $\frac{ab^4c^2}{8}$

## Solutions to trial exercise problems

3.  $5a^0 = 5 \cdot 1 = 5$  10.  $4z^{-2} = 4 \cdot \frac{1}{z^2} = \frac{4}{z^2}$   
 15.  $2x^{-4}y^2 = 2 \cdot \frac{1}{x^4} \cdot y^2 = \frac{2y^2}{x^4}$  32.  $\frac{3^4}{3^2} = 3^{4-2} = 3^2 = 9$   
 36.  $\frac{x^4x^3}{x^2} = \frac{x^{4+3}}{x^2} = \frac{x^7}{x^2} = x^{7-2} = x^5$  48.  $\frac{5^2a^3b}{5^3a^7b^3}$   
 $= 5^{2-3}a^{3-7}b^{1-3} = 5^{-1}a^{-4}b^{-2} = \frac{1}{5^1} \cdot \frac{1}{a^4} \cdot \frac{1}{b^2} = \frac{1}{5a^4b^2}$   
 49.  $x^{-4}x^7 = x^{-4+7} = x^3$  Alternate:  $x^{-4}x^7 = \frac{1}{x^4} \cdot x^7 = \frac{x^7}{x^4} = x^{7-4}$   
 $= x^3$  57.  $(-5)^{-3} = \frac{1}{(-5)^3} = -\frac{1}{125}$  (Note: The sign of the base,  
 $-5$ , is unchanged.) 73.  $\frac{4^{-1}a^{-2}b^3c^0}{2a^{-3}b^{-1}c^{-2}} = \frac{\frac{1}{4^1} \cdot \frac{1}{a^2} \cdot b^3 \cdot 1}{2 \cdot \frac{1}{a^3} \cdot \frac{1}{b^1} \cdot \frac{1}{c^2}} = \frac{\frac{b^3}{4a^2}}{\frac{2}{a^3bc^2}}$   
 $= \frac{b^3}{4a^2} \cdot \frac{a^3bc^2}{2} = \frac{a^3b^3 + 1c^2}{2 \cdot 4a^2} = \frac{a^3b^4c^2}{8a^2} = \frac{b^4c^2}{8} \cdot a^{3-2} = \frac{b^4c^2}{8} \cdot a^1$   
 $= \frac{ab^4c^2}{8}$

## Review exercises

1.  $-10$  2.  $21$  3.  $4$  4.  $-16$  5.  $a^8$   
 6.  $x^{12}$  7.  $x^6$  8.  $4a^2b^2$

## Exercise 3-4

## Answers to odd-numbered problems

1.  $8a^6$  3.  $8x^6y^3$  5.  $x^{16}y^{12}z^4$  7.  $125a^{15}b^6c^3$   
 9.  $\frac{1}{4a^4}$  11.  $\frac{16}{x^4}$  13.  $\frac{y^{12}}{27x^3}$  15.  $\frac{y^{10}}{x^4z^6}$   
 17.  $6x^7y^3$  19.  $-6x^{10}y^4$  21.  $6x^6y^3z^7$  23.  $\frac{8x^3}{y^6}$   
 25.  $\frac{9a^4}{b^6}$  27.  $\frac{8x^6}{y^9}$  29.  $\frac{4x^6y^6}{z^{10}}$  31.  $\frac{b^8}{a^5}$   
 33.  $\frac{R^2}{3S^4}$  35.  $\frac{4}{ab^9}$  37.  $\frac{3S^3}{R^4}$  39.  $a^{10}b^{17}$   
 41.  $32a^{12}$  43.  $32x^{14}y^{13}$  45.  $\frac{y^2}{xz^4}$   
 47.  $\frac{a^6b^{10}}{4}$  49.  $\frac{b^6}{a^3c^3}$

## Solutions to trial exercise problems

9.  $(2a^2)^{-2} = \frac{1}{(2a^2)^2} = \frac{1}{2^2(a^2)^2} = \frac{1}{2^2a^4} = \frac{1}{4a^4}$   
 31.  $\frac{a^{-2}b^3}{a^3b^{-5}} = \frac{\frac{1}{a^2} \cdot b^3}{a^3 \cdot \frac{1}{b^5}} = \frac{\frac{b^3}{a^2}}{\frac{a^3}{b^5}} = \frac{b^3}{a^2} \cdot \frac{b^5}{a^3} = \frac{b^{3+5}}{a^{2+3}} = \frac{b^8}{a^5}$   
 39.  $(a^2b^3)^3(ab^2)^4 = (a^2)^3(b^3)^3 \cdot a^4(b^2)^4 = a^6b^9a^4b^8 = a^{6+4}b^{9+8} = a^{10}b^{17}$

## Review exercises

1.  $35.34$  2.  $10.36$  3.  $16.72$  4.  $28.98$  5.  $18.81$  6.  $49.5$



## Exercise 3–5

## Answers to odd-numbered problems

1.  $2.55 \times 10^2$  3.  $1.2345 \times 10^4$  5.  $1.55 \times 10^5$   
 7.  $8.55076 \times 10^2$  9.  $1.0076 \times 10^6$  11.  $1.2 \times 10^{-4}$   
 13.  $8.1 \times 10^{-6}$  15.  $7 \times 10^{-4}$  17.  $9.4 \times 10^{-11}$   
 19.  $-4.5 \times 10^3$  21.  $-5.85 \times 10^6$  23.  $-4.578 \times 10$   
 25.  $-2.985 \times 10^{-8}$  27. 49,900,000 29. 7.23 31. 0.0042  
 33. 0.00000147 35. 0.000789 37.  $-0.00000000482$   
 39.  $-0.00000492$  41.  $1 \times 10^{-9}$  43.  $2 \times 10^{12}$   
 45. 35,600,000 47. 0.000 000 000 000 000 000 093  
 49. 140,000 51.  $1.22304 \times 10^{14}$  53.  $3.63226 \times 10^{-1}$   
 55.  $1.76979 \times 10^{-7}$  57.  $4.84481 \times 10^8$  59.  $1.4 \times 10^3$   
 61.  $4.6 \times 10^3$

## Solution to trial exercise problem

$$\begin{aligned} 58. (177,000) \div (0.15) &= \frac{1.77 \times 10^5}{1.5 \times 10^{-1}} \\ &= \frac{1.77}{1.5} \times 10^{5-(-1)} = \frac{1.77}{1.5} \times 10^6 = 1.18 \times 10^6 \end{aligned}$$

## Review exercises

1.  $5x$  2.  $3a^2$  3.  $12ab$  4.  $x^3 + 2x^2$  5.  $6a^2 - 15a$   
 6.  $3x^3y + 2x^2y^2 - 7x^2y$

## Chapter 3 review

1.  $a^{12}$  2.  $a^{14}$  3.  $4^5 = 1,024$  4.  $x^4y^4$  5.  $a^{15}$  6.  $20a^4b^5$   
 7.  $6x^3y^7$  8.  $-15x^5$  9.  $6a^3b^5$  10.  $10x^5y^5$  11.  $-6a^6b^{10}$   
 12.  $15x^2 - 10xy$  13.  $-6a^4b + 9a^3b^2 - 12a^2b^3$   
 14.  $30x^2y^2 - 10xy^3$  15.  $x^2 - x - 12$  16.  $x^2 + 10x + 25$   
 17.  $a^2 - 49$  18.  $15x^2 + 7xy - 2y^2$  19.  $x^3 + x^2y - 5xy^2 - 2y^3$   
 20.  $3a^3 - 4a^2b - 5ab^2 + 2b^3$  21.  $8a^3 + 12a^2b + 6ab^2 + b^3$   
 22.  $\frac{1}{b^2}$  23.  $\frac{5}{a^2}$  24.  $a^4$  25.  $\frac{1}{x^3}$  26. 1 27.  $\frac{5}{x^3y^2}$  28.  $a^3$   
 29.  $\frac{a^5}{b^5}$  30.  $\frac{4y^2z^2}{x^2}$  31.  $\frac{b^3}{4a^3}$  32.  $\frac{a^9}{b^3}$  33.  $8a^6b^9$   
 34.  $3^{12}x^{16}y^{20} = 531,441x^{16}y^{20}$  35.  $\frac{y^6}{4x^2}$  36.  $\frac{xy^3z^3}{2}$  37.  $\frac{2a^2c^3}{b^6}$   
 38.  $8x^{26}y^{25}$  39.  $72a^{14}b^3$  40.  $\frac{a^9b^3}{c^{12}}$  41.  $\frac{9a^6b^4}{c^{10}}$  42.  $\frac{32x^5y^{20}}{z^{30}}$   
 43.  $1.84 \times 10^3$  44.  $1.57 \times 10^{-3}$  45.  $1.07 \times 10^8$   
 46.  $8.49 \times 10^{11}$  47.  $-3.75 \times 10$  48.  $-5.43 \times 10^{-3}$   
 49. 504,000 50. 0.00639 51.  $-596$  52.  $-0.00886$   
 53. 0.000000735 54. 812,000,000 55.  $2.67672 \times 10^5$   
 56.  $1.54818 \times 10^{-8}$  57.  $7.2 \times 10^{-3}$  58.  $1.25 \times 10^{-7}$

## Chapter 3 cumulative test

1. false 2. false 3. true 4.  $-2$  5. undefined 6. 8  
 7. 0 8. 16 9. 38 10.  $x^2y + 2xy^2 - 3x^2y^2$  11.  $-40$   
 12. 0 13.  $9x^2 - 6xy + y^2$  14. 0 15.  $4a^4b^{10}$  16.  $-25$   
 17. 0 18.  $9x^2 - 4y^2$  19.  $6x^3y^5$  20. 26 21. undefined  
 22.  $4a + 2b$  23.  $x^3 - 2x - 1$  24.  $x^2$  25.  $a^4$  26.  $\{7\}$   
 27.  $\left\{\frac{5}{3}\right\}$  28.  $\left\{-\frac{19}{4}\right\}$  29.  $x > -\frac{1}{3}$  30.  $x > \frac{1}{6}$   
 31.  $\left\{-\frac{3}{2}\right\}$  32.  $-8 \leq x \leq -1$  33. 97 34. 72  
 35. \$18,500 at 8%; \$11,500 at 7%

## Chapter 4

## Exercise 4–1

## Answers to odd-numbered problems

1.  $2(y + 3)$  3.  $4(x^2 + 2y)$  5.  $3(x^2y + 5z)$   
 7.  $7(a - 2b + 3c)$  9.  $3(5xy - 6z + x^2)$   
 11.  $7(6xy - 3y^2 + 1)$  13.  $2(4x - 5y + 6z - 9w)$   
 15.  $5ab(4a - 12 + 9b)$  17.  $3xy(x + 2)$  19.  $2R^2(R^2 - 3)$   
 21.  $x(2x^2 - x + 1)$  23.  $3ab(5 + 6b - a)$  25.  $xy(y + z + 1)$   
 27.  $2L(L^2 - 9 + L)$  29.  $5p(p + 2 + 3p^2)$  31.  $-3(2x + 3)$   
 33.  $2(3x - 4z - 6w)$  35.  $-3(4L - 5W + 2H)$   
 37.  $-x(1 - x + x^2)$  39.  $-xyz(1 - x + y - z)$   
 41.  $-5ab(2ab - 3 + 4a^2b^2)$  43.  $(a + b)(x + y)$   
 45.  $5(2a + b)(3x + 2y)$  47.  $3(a + 4b)(x + 2y)$   
 49.  $(b + 6)(8a - 1)$  51.  $(a + b)(c + d)$   
 53.  $(2a + b)(3x - 2y)$  55.  $(2x + y)(2a - b)$   
 57.  $(5x - 3y)(4x + z)$  59.  $(a + 3b)(4x - 3y)$   
 61.  $(2c - y)(a + 3b)$  63.  $(c + 4y)(2a + 3b)$   
 65.  $(3x + y)(2a + b)$  67.  $(x - 2d)(3a + b)$   
 69.  $(a + 5)(2a^2 + 3)$  71.  $(4a^2 + 3)(2a - 1)$

73.  $\pi r(s + r)$  75.  $\frac{wx}{48EI}(2x^3 - 3lx^2 - l^3)$

## Solutions to trial exercise problems

11.  $42xy - 21y^2 + 7 = 7 \cdot 6xy - 7 \cdot 3y^2 + 7 \cdot 1$   
 $= 7(6xy - 3y^2 + 1)$  25.  $xy^2 + xyz + xy = xy \cdot y + xy \cdot z$   
 $+ xy \cdot 1 = xy(y + z + 1)$  33.  $6x - 8z - 12w = 2(\quad)$ ;  
 $2 \cdot 3x - 2 \cdot 4z - 2 \cdot 6w = 2(3x - 4z - 6w)$   
 34.  $-4a^3 - 36ab + 16ab^2 - 24b^3 = -4(\quad)$ ;  
 $(-4)a^3 + (-4)(9ab) + (-4)(-4ab^2) + (-4)(6b^3)$   
 $= -4(a^3 + 9ab - 4ab^2 + 6b^3)$  36.  $-3a + a^3b = -a(\quad)$ ;  
 $(-a) \cdot 3 + (-a)(-a^2b) = -a(3 - a^2b)$  45.  $15x(2a + b)$   
 $+ 10y(2a + b) = 5(2a + b) \cdot 3x + 5(2a + b) \cdot 2y$   
 $= 5(2a + b)(3x + 2y)$  53.  $6ax - 2by + 3bx - 4ay$   
 $= 6ax + 3bx - 4ay - 2by = (6ax + 3bx) + (-4ay - 2by)$   
 $= 3x(2a + b) - 2y(2a + b) = (2a + b)(3x - 2y)$   
 75.  $Y = \frac{2wx^4}{48EI} - \frac{3lwx^3}{48EI} - \frac{l^3wx}{48EI} = \frac{wx}{48EI} \cdot 2x^3 - \frac{wx}{48EI} \cdot 3lx^2$   
 $- \frac{wx}{48EI} \cdot l^3 = \frac{wx}{48EI}(2x^3 - 3lx^2 - l^3)$

## Review exercises

1. 4, 5 2. 3, 4 3.  $-8, 2$  4. 8,  $-2$  5. 8, 2 6.  $-8, -2$   
 7. 6, 6 8. 11, 1

## Exercise 4–2

## Answers to odd-numbered problems

1.  $(a + 6)(a + 3)$  3.  $(x + 12)(x - 1)$  5.  $(y + 15)(y - 2)$   
 7.  $(x - 12)(x - 2)$  9.  $(a + 8)(a - 3)$  11.  $(x + 6)(x + 2)$   
 13.  $(a - 6)(a + 4)$  15.  $2(x + 5)(x - 2)$  17.  $3(x - 8)(x + 2)$   
 19. will not factor, prime polynomial 21.  $(y + 15)(y + 2)$   
 23.  $4(x + 2)(x - 3)$  25.  $5(a - 5)(a + 2)$   
 27.  $(xy - 6)(xy + 3)$  29.  $(xy + 12)(xy + 1)$   
 31.  $3(xy + 3)(xy - 4)$  33.  $(x + 2y)(x + y)$   
 35.  $(a - 3b)(a + b)$  37.  $(a - 3b)(a + 2b)$   
 39.  $(x + 3y)(x - 5y)$

# Campfire queen Cycling champion Sentimental geologist\*

Learn more about  
Marjon Walrod  
and tell us more  
about you. Visit  
[pwc.com/bringit](http://pwc.com/bringit).

Your life. You can  
bring it with you.



\*connectedthinking

PRICewaterhouseCOOPERS 

© 2006 PricewaterhouseCoopers LLP. All rights reserved. "PricewaterhouseCoopers" refers to PricewaterhouseCoopers LLP (a Delaware limited liability partnership) or, as the context requires, the PricewaterhouseCoopers global network or other member firms of the network, each of which is a separate and independent legal entity. \*connectedthinking is a trademark of PricewaterhouseCoopers LLP (US). We are proud to be an Affirmative Action and Equal Opportunity Employer.



# Contents

20 Point Learning System	xi
Preface	xvii
Study Tips	xxiii

## Chapter 1 ■ Operations with real numbers



1-1 Operations with fractions	2
1-2 Operations with decimals and percents	14
1-3 The set of real numbers and the real number line	25
1-4 Addition of real numbers	34
1-5 Subtraction of real numbers	41
1-6 Multiplication of real numbers	46
1-7 Division of real numbers	51
1-8 Properties of real numbers and order of operations	56
Chapter 1 lead-in problem	62
Chapter 1 summary	63
Chapter 1 error analysis	64
Chapter 1 critical thinking	64
Chapter 1 review	64

## Chapter 2 ■ Solving equations and inequalities




2-1 Algebraic notation and terminology	67
2-2 Evaluating algebraic expressions	72
2-3 Algebraic addition and subtraction	79
2-4 The addition and subtraction property of equality	86
2-5 The multiplication and division property of equality	93
2-6 Solving linear equations	98
2-7 Solving literal equations and formulas	104
2-8 Word problems	107
2-9 Solving linear inequalities	113
Chapter 2 lead-in problem	122
Chapter 2 summary	122
Chapter 2 error analysis	123
Chapter 2 critical thinking	123
Chapter 2 review	124
Chapter 2 cumulative test	125



**Chapter 3 ■ Polynomials and exponents**

3-1	Exponents—I	127
3-2	Products of algebraic expressions	133
3-3	Exponents—II	139
3-4	Exponents—III	145
3-5	Scientific notation	148
	Chapter 3 lead-in problem	151
	Chapter 3 summary	152
	Chapter 3 error analysis	152
	Chapter 3 critical thinking	152
	Chapter 3 review	153
	Chapter 3 cumulative test	154

**Chapter 4 ■ Factoring and solution of quadratic equations by factoring**

4-1	Common factors	155
4-2	Factoring trinomials of the form $x^2 + bx + c$	162
4-3	Factoring trinomials of the form $ax^2 + bx + c$	166
4-4	Factoring the difference of two squares and perfect square trinomials	175
4-5	Other types of factoring	179
4-6	Factoring: A general strategy	184
4-7	Solving quadratic equations by factoring	186
4-8	Applications of the quadratic equation	193
	Chapter 4 lead-in problem	197
	Chapter 4 summary	198
	Chapter 4 error analysis	198
	Chapter 4 critical thinking	198
	Chapter 4 review	199
	Chapter 4 cumulative test	200

**Chapter 5 ■ Rational Expressions, Ratio and Proportion**

5-1	Rational numbers and rational expressions	202
5-2	Simplifying rational expressions	207
5-3	The quotient of two polynomials	212
5-4	Ratio and proportion	219
	Chapter 5 lead-in problem	228
	Chapter 5 summary	228
	Chapter 5 error analysis	228
	Chapter 5 critical thinking	229
	Chapter 5 review	229
	Chapter 5 cumulative test	230



## Chapter 6 ■ Operations with Rational Expressions



6-1	Multiplication and division of rational expressions	232
6-2	Addition and subtraction of rational expressions	239
6-3	Addition and subtraction of rational expressions	245
6-4	Complex fractions	253
6-5	Rational equations	258
6-6	Rational expression applications	265
	Chapter 6 lead-in problem	270
	Chapter 6 summary	271
	Chapter 6 error analysis	271
	Chapter 6 critical thinking	272
	Chapter 6 review	272
	Chapter 6 cumulative test	274

## Chapter 7 ■ Linear Equations in Two Variables




7-1	Ordered pairs and the rectangular coordinate system	276
7-2	Graphs of linear equations	289
7-3	The slope of a line	297
7-4	The equation of a line	305
7-5	Graphing linear inequalities in two variables	315
7-6	Functions defined by linear equations in two variables	323
	Chapter 7 lead-in problem	329
	Chapter 7 summary	329
	Chapter 7 error analysis	330
	Chapter 7 critical thinking	330
	Chapter 7 review	331
	Chapter 7 cumulative test	333

## Chapter 8 ■ Systems of Linear Equations




8-1	Solutions of systems of linear equations by graphing	335
8-2	Solutions of systems of linear equations by elimination	340
8-3	Solutions of systems of linear equations by substitution	346
8-4	Applications of systems of linear equations	351
8-5	Solving systems of linear inequalities by graphing	360
	Chapter 8 lead-in problem	363
	Chapter 8 summary	364
	Chapter 8 error analysis	364
	Chapter 8 critical thinking	364
	Chapter 8 review	365
	Chapter 8 cumulative test	366

## Chapter 9 ■ Roots and Radicals



9-1	Principal roots	367
9-2	Product property for radicals	373
9-3	Quotient property for radicals	377
9-4	Sums and differences of radicals	383
9-5	Further operations with radicals	386
9-6	Fractional exponents	391
9-7	Equations involving radicals	395
	Chapter 9 lead-in problem	400
	Chapter 9 summary	400
	Chapter 9 error analysis	401
	Chapter 9 critical thinking	401
	Chapter 9 review	401
	Chapter 9 cumulative test	402

## Chapter 10 ■ Solutions of Quadratic Equations



10-1	Solutions of quadratic equations by extracting the roots	404
10-2	Solutions of quadratic equations by completing the square	409
10-3	Solutions of quadratic equations by the quadratic formula	416
10-4	Complex solutions to quadratic equations	424
10-5	The graphs of quadratic equations in two variables—quadratic functions	432
	Chapter 10 lead-in problem	444
	Chapter 10 summary	445
	Chapter 10 error analysis	445
	Chapter 10 critical thinking	446
	Chapter 10 review	446
	Final examination	447
	Appendix Answers and Solutions	449
	Index	505



# Index

## A

Abscissa of a point, 281  
Absolute value, 31  
Addition  
  of decimals, 16  
  of fractions, 7–9, 239  
  of rational expressions, 239, 246  
Addition, identity element of, 35  
Addition, properties of  
  associative, 38  
  commutative, 35  
  identity, 35  
  rational expressions, 239, 246  
Addition and subtraction property of equality, 88, 340  
Addition and subtraction property of inequalities, 115  
Addition of  
  algebraic expressions, 80–81  
  like radicals, 384–85  
  more than two real numbers, 42  
  rational expressions, 239, 246  
  two negative numbers, 36  
  two numbers with different signs, 36–37  
  two positive numbers, 34  
  two real numbers, 38  
Additive inverse, 37–38  
Algebraic expression, 67  
Approximately equal to, 19, 29, 369  
Associative property of  
  addition, 38  
  multiplication, 49  
Axes, 280  
Axis of symmetry, 437

## B

Base, 56, 127  
Binomial, 68  
Binomial, square of, 135, 410  
Boundary line, 316  
Braces, 25, 42  
Brackets, 42

## C

Cantor, Georg, 25  
Coefficient, 68  
Common denominator, 7  
Commutative property of  
  addition, 35  
  multiplication, 46  
Completely factored form, 157

Completing the square, 411–13  
Complex fractions, 253  
  primary denominator, 253  
  primary numerator, 253  
  secondary denominators of, 253  
  simplifying, 253–55  
Complex numbers, 425  
  addition of, 425  
  multiplication of, 426  
  rationalizing the denominator of, 427  
Components of an ordered pair, 277  
Compound inequality, 114  
Conjugate factors, 388  
Consistent system of equations, 337  
Constant, 67  
Coordinate, 29, 281  
Coordinates of a point, 281  
Counting numbers, 26

## D

Decimal number, 14  
  fraction, 14  
  point, 14  
  repeating, 19, 27  
  terminating, 27  
Decreasing, 29  
Degree of a polynomial, 69, 186  
Denominator  
  of a fraction, 2  
  least common, 42  
  of a rational expression, 203  
Denominator, rationalizing a, 427, 480, 482  
Dependent system, 337  
Descending powers, 69, 163  
Difference of two squares, 136  
Discriminant, 429  
Distance-rate-time, 266, 352  
Distributive property of  
  multiplication over addition, 79  
Dividend, 51  
Division  
  monomial by monomial, 212  
  polynomial by monomial, 212  
  polynomial by polynomial, 213–16  
  property of rational expressions, 234  
  of rational expressions, 235  
Division, definition of, 51  
Division by zero, 53  
Division involving zero, 53  
Division of  
  decimals, 17  
  fractions, 5, 235  
  like bases, 140  
  two or more real numbers, 52  
  two real numbers, 52

Divisor, 51  
Domain  
  of a rational expression, 204–5  
  of a relation, 323

## E

Element of a set, 25  
Elimination, method of solving systems of  
  linear equations, 340–44  
Empty set, 260, 398  
Equality, properties of  
  addition and subtraction, 88  
  multiplication and division, 94  
  squaring, 396  
  symmetric, 89  
Equation, parts of an, 86  
Equation of a line, 305  
Equations  
  conditional, 87  
  diagram of, 86  
  equivalent, 87  
  first-degree, 87  
  graph of, 282–83, 289–94  
  identical, 87  
  linear, 87, 276  
  literal, 104  
  quadratic, 186, 404–31  
  radical, 395–96  
  rational, 258  
  system of, 335  
Equivalent rational expressions, 245–46  
Evaluating rational expressions, 203–4  
Evaluation, 72  
Expanded form, 56, 127  
Exponent, 56, 127  
Exponential form, 56, 127  
Exponents, properties and definitions of  
  definition, 127  
  fraction, 139  
  fractional, 392  
  group of factors to a power, 129  
  negative exponents, 141  
  power of a power, 130  
  product, 128  
  quotient, 140  
  zero as an exponent, 142  
Extracting the roots, 405–6  
Extraneous solutions, 260, 396

## F

Factor, 3, 46  
Factor, greatest common, 156  
Factored form, completely, 157

**Factoring**

- common factors, 156
- difference of two cubes, 179–80
- difference of two squares 175–76
- four-term polynomials, 159
- by inspection, 170–74
- perfect square trinomials, 177–78
- strategy, 184
- sum of two cubes, 181–82
- trinomials, 162–64, 166–74

**FOIL, 135****Formulas, 74, 104****Four-term polynomials, 159****Fraction, 2**

- complex, 253
- improper, 2
- proper, 2

**Fraction exponents, 392****Fraction to a power, 139****Function, 324**

- domain of a, 325
- range of a, 325

**Fundamental principle of rational expressions, 207****G****Graph**

- of a linear equation, in two variables, 282–83, 289–94
- of a point, 29, 359
- a quadratic equation, 432–40
- of systems of equations, 336–37

**Graphing**

- a linear equation in two variables, 289–94
- linear inequalities in two variables, 315–19
- quadratic functions, 434–39
- systems of linear equations, 336–37
- systems of linear inequalities, 360–61

**Greatest common factor, 155–56****Grouping symbols, 42, 82****Grouping symbols, removing, 82****Group of factors to a power, 129****H****Half-planes, 316****Horizontal line, 292–93**

- slope of, 300

**I****Identical equation, 87****Identity, 87****Identity element**

- of addition, 35
- of multiplication, 47

**Inconsistent system, 337****Increasing, 29****Independent system, 337****Indeterminate, 53****Index, 370****Inequalities**

- compound, 114

**linear, 113**

- strict, 30
- in two variables, 315–19
- weak, 30

**Inequalities, properties of**

- addition and subtraction, 115
- multiplication and division, 115

**Inequality symbols, 30, 113****Integers, 26****Intercepts, x- and y-, 290****Inverse property**

- additive, 38
- multiplicative, 94

**Irrational numbers, 27, 368****L****Least common denominator (LCD), 7, 242****Least common multiple (LCM), 242****Like bases, 128****Like radicals, 383****Like terms, 80****Linear equation, 87****Linear equations in two variables, 276**

- graphs of, 289–93
- systems of, 335

**Linear inequalities in two variables, 315**

- graphs of, 315–19

**Linear inequality, 113****Lines**

- equation of, 305
- horizontal, 300
- parallel, 309
- perpendicular, 310
- point-slope form, 306
- slope-intercept form of, 307
- slope of, 297–302
- standard form of, 305
- vertical, 301

**Literal equation, 104****M****Mathematical statement, 86****Member of an equation, 86****Member of a set, 25****Mixed number, 5****Monomial, 68****Multinomial, 69****Multiple, least common, 242****Multiplication**

- of decimals, 16
- of fractions, 4
- identity element of, 47
- of rational expressions, 233
- symbols for, 46

**Multiplication, properties of**

- associative, 49
- commutative, 46
- fractions, 4, 232
- identity, 47

- rational expressions, 233

**Multiplication and division property for inequalities, 115****Multiplication and division property of equality, 94****Multiplication of**

- like bases, 128
- monomial and a multinomial, 133
- monomials, 130
- multinomials, 134
- $n$ th roots, 374
- square roots, 373
- two negative numbers, 47
- two numbers with different signs, 47
- two or more real numbers, 48
- two positive numbers, 46
- two real numbers, 48

**Multiplicative inverse, 94****N****Natural numbers, 26****Negative exponents, 141****Negative integers, 26****Negative of, 31** **$n$ th root**

- addition of, 383–85
- definition of, 370
- of a fraction, 378, 380
- rationalizing, 379–81, 388

**Number line, 29****Numerator, 2, 203****Numerical coefficient, 68****O****Open sentence, 86****Opposite of, 31****Ordered pairs, first component of, 277****Ordered pairs, second component of, 277****Ordered pairs of numbers, 277****Order of operations, 57****Order relationships, 30****Ordinate of a point, 281****Origin, 29, 280****of a rectangular coordinate plane, 280****P****Parabola, 434**

- axis of symmetry of a, 437
- graphing a, 434–39
- intercepts of a, 435
- vertex of a, 436–37

**Parallel lines, 309****Parentheses, 42****Percent, 19–20****Percentage, 20–21****Perfect cube, 180****Perfect square, 175–76****Perfect square integer, 368****Perfect square trinomial, 135, 409****Perimeter, 10, 74****Perpendicular lines, 310****Pi ( $\pi$ ), 28****Point**

- abscissa of a, 281
- graph of a, 281
- ordinate of a, 281

**Point-slope form, 306****Polynomial, 68**



Polynomial, degree of, 69, 186  
 Positive integers, 26  
 Power of a power, 130  
 Prime factor form, 3  
 Prime number, 3  
 Prime polynomial, 164  
 Principal  $n$ th root, 370  
 Principal square root, 368  
 Product, 3, 46  
 Product of rational expressions, 233  
 Product property of exponents, 128  
 Product property of radicals, 374  
 Product property of square roots, 373  
 Properties of real numbers, 56  
 Proportion, 221  
   property of, 221  
   terms of, 221  
 Pythagorean Theorem, 369

**Q**

Quadrants, 280  
 Quadratic  
   equations, 186, 404–31  
   formula, 417–19  
   standard form of, 187, 404  
 Quotient, 5, 51  
   of two polynomials, 212–16  
 Quotient property of exponents, 140  
 Quotient property of radicals, 378, 380

**R**

Radical equation, 395–96  
 Radicals, properties of  
   product, 373, 374  
   quotient, 378, 380  
 Radical symbol, 368  
 Radicand, 368  
 Range of a relation, 323  
 Ratio, 219  
 Rational equation, 258  
   application, 265–67  
   in more than one variable, 261  
 Rational expression, 202  
 Rational expressions  
   applications of, 265–67  
   completely reduced, 208–10  
   definition of, 202  
   denominator of a, 203  
   difference of, 239, 246  
   domain of, 204  
   fundamental principle of, 207  
   least common denominator of, 242  
   numerator of, 203  
   product of, 233  
   quotient of, 235  
   sum of, 239, 246  
 Rationalizing a denominator, 379–81, 388, 427  
 Rational number, 27  
 Real number, 28  
 Real number line, 29  
 Reciprocal, 5, 94  
 Rectangular coordinate plane, 280  
 Reducing to lowest terms, 4, 208

Relation, 323  
   domain of a, 323  
   range of a, 323  
 Remainder, 5  
 Repeating decimal, 19  
 Root, 86  
 Root, extraneous, 260

**S**

Scientific notation, 148  
   computation using, 149  
   standard form of, 149  
 Set, 25  
 Set symbolism, 25  
 Signed numbers, 34  
 Similar terms, 80  
 Simple interest, 112  
 Simplifying rational expressions, 207  
 Slope, 297–98  
   definition of, 298–99  
   of a horizontal line, 300  
   of parallel lines, 309  
   of perpendicular lines, 310  
   undefined, 300–1  
   of a vertical line, 301  
 Slope-intercept form, 307  
 Solution, 86, 276  
   extraneous, 260  
 Solution set, 86, 187  
   of a linear equation in two variables, 277  
 Solutions of quadratic equations  
   by completing the square, 411–13  
   complex solutions of, 427  
   by extracting the roots, 405–6  
   by factoring, 404–5  
   by quadratic formula, 418–19  
 Special products, 135–37  
 Square of a binomial, 135  
 Square root, 367  
 Square root property, 405  
 Standard form of a quadratic equation, 404  
 Standard form of the equation of a line, 305  
 Statement, mathematical, 86  
 Straight line, 289  
 Strict inequality, 30  
 Subscripts, 75  
 Subset, 26  
 Substitution property, 72  
   method of solving a system of linear equations, 346–48  
 Subtraction, definition of, 41  
   decimals, 16  
 Subtraction, of rational expressions, 239  
 Subtraction of  
   fractions, 7–9, 239  
   more than two real numbers, 42  
   two real numbers, 41  
 Symbols  
   absolute value, 31  
   braces, 25, 42  
   brackets, 42  
   is an element of, 26  
   is approximately equal to, 19  
   is a subset of, 26  
   is greater than, 30  
   is greater than or equal to, 30

is less than, 30  
 is less than or equal to, 30  
 multiplication dot, 46  
 null set, 260  
 parentheses, 42  
 Symmetric property of equality, 89  
 Symmetry, 31  
 Systems of linear equations, 335  
   applications of, 351–55  
   consistent and independent, 337–38  
   dependent, 337–38  
   graphing of, 336–37  
   inconsistent, 337–38  
   linear, 335  
   solution by elimination, 340–44  
   solution by substitution, 346–48  
 Systems of linear inequalities, 360–61

**T**

Term, 67  
 Trinomial, 68  
 Trinomial, perfect square, 135, 409

**U**

Undefined, 53  
 Undefined slope, 301  
 Undirected distance, 31

**V**

Variable, 29, 67  
 Vertex of a parabola, 436–37  
 Vertical line, 293  
   slope of, 301

**W**

Weak inequality, 30  
 Whole numbers, 26

**X**

$x$ -axis, 280  
 $x$ -intercept, 290, 435

**Y**

$y$ -axis, 280  
 $y$ -intercept, 290, 435

**Z**

Zero, division involving, 53  
 Zero as an exponent, 142  
 Zero factor property, 47  
 Zero product property, 187